

The Relative Distribution of Otoliths as a Means of Larval Fish Identification

Li Shu Chen^{1,*} and Hong Young Yan²

¹National Museum of Marine Science and Technology – Preparatory Office, Keelung, Taiwan 202, R.O.C.

²School of Biological Sciences, University of Kentucky, Lexington, KY 40506-0225, USA

(Accepted December 21, 2001)

Li Shu Chen and Hong Young Yan (2001) The relative distribution of otoliths as a means of larval fish identification. *Zoological Studies* 41(2): 144-152. A recently developed auditory brainstem response (ABR) protocol has indicated that acoustically evoked brainwaves in fishes are species-specific. These brainwaves reflect the spatial distribution of auditory nuclei and the 3 auditory end organs along the ascending auditory neuronal pathways. Species-specific brainwaves suggest the hypothesis that the distribution of otoliths, which are closely coupled with sensory hair cells whose evoked response forms the 1st wave of the ABR, should also be species-specific. Eight species of marine fish larvae (belonging to 8 genera and 4 families) collected from Taiwan and Texas, USA were cleared in trypsin and examined microscopically with double polarization filters. Three indices (inter-utricular ratio, IUR; inter-sacculus ratio, ISR; and inter-lagena ratio, ILR) were developed to measure the ratio of inter-otolith to inter-orbital distances. IUR and ISR succeeded in separating species, but ILR was not as useful because of late development of the lagena. This study supports the hypothesis that the distribution of otoliths in fish larvae differs among species.
<http://www.sinica.edu.tw/zool/zoolstud/41.2/144.pdf>

Key words: Mechanosensory physiology, Inner ear, Otolith, Auditory pathway, Taxonomy.

The vestibular and auditory sensory modalities of fish are encoded by 3 major otolithic endorgans: the utricle, the saccule, and the lagena. Within each otic chamber resides an otolith: the lapillus (utricle chamber), the sagitta (saccule chamber), and the astericus (lagena chamber). Each otolith is lined with patches of sensory epithelia composed of sensory hair cells and supporting cells. An otolithic membrane provides a mechanical linkage between the otolith and the ciliary bundles of sensory hair cells where acoustic and vestibular stimuli are encoded. The otoliths of adult fish are calcareous structures of characteristic shapes and sizes and are species-specific (Helfman et al. 1997). One of the otoliths in particular, the sagitta, has been used as a major character in taxonomic studies of adult salmon (Casteel 1974), sciaenids (Chao 1978), neopterygian marine fishes (Hecht 1978), and fossil fishes (Huygeraert and Nolf 1979). Since the discovery of daily growth rings in otoliths of lar-

val fish (Pannella 1971 1980), great interest has been focused on age, growth, mortality, life histories, migration, recruitment, and settlement of larvae from the information retrieved from the larval otoliths of American and European eels (*Anguilla rostrata*, *A. anguilla*) (Wang and Tzeng 2000), anchovies (*Encrasicholina punctifer*, *Engraulis japonicus*) (Wang and Tzeng 1999), Atlantic herring (*Clupea harengus*) (Townsend and Graham 1981, Lough et al. 1982), Atlantic mackerel (*Scomber scombrus*) (Kendall and Gordon 1981), burbot (*Lota lota*) (Fischer 1999), chinook salmon (*Oncorhynchus tshawytscha*) (Zhang and Beamish 2000), grey mullet (*Mugil cephalus*) (Chang et al. 2000), haddock (*Melanogrammus aeglefinus*) (Gallego et al. 1999), Japanese eel (*Anguilla japonica*) (Tzeng and Tsai 1994, Tzeng et al. 1994, Tzeng 1995a, b, Cheng and Tzeng 1996), nehu (*Stolephorus purpureus*) (Struhsaker and Uchiyama 1976), northern anchovy (*Engraulis mordax*) (Methot and Kramer 1979), Pa-

*To whom correspondence and reprint requests should be addressed. Tel: 886-2-24631841 ext. 301. Fax: 886-2-24627716. E-mail: Lschen@mail.ntou.edu.tw

cific hake (*Merluccius productus*) (Bailey 1982), pleuronectid (*Parophrys vetulus*) (Laroche et al. 1982), red drum (*Sciaenops ocellatus*) (Rooker et al. 1999), rockfish (*Sebastes thompsoni*) (Kokita and Omori 1999), weak fish (*Cynoscion regalis*) (Thorrold et al. 1998, Paperno et al. 2000) and other marine (Brothers et al. 1976 1983, Lou and Moltschaniwskyj 1992, Wilson and McCormick 1997 1999) as well as freshwater fishes (Claramunt and Wahl 2000).

However, to date, otoliths have rarely been used for larval species identification. Constraints of the limited use of otolith for species identification are: 1) small size, 2) simpler form, and 3) apparent lack of obvious distinguishing external features (Brothers 1984). Some progress has been made by examining crystallographic, mineralogical, and chemical levels (Lowenstan and Fitch 1978 1981, Lowenstan 1980 1981), amino acid composition (Degens et al. 1969), and trace elements analysis (Papadopoulou et al. 1978 1980, Gaudie et al. 1980) of adult otoliths as potential characters for taxonomic use. The direct application of these techniques to larval fish, however, is largely hampered by the small size of larval otoliths which renders analyses difficult or next to impossible.

Recently an auditory brainstem response (ABR) recording technique was developed to investigate acoustically evoked brainwaves from ascending auditory neuronal pathways of fishes (Kenyon et al. 1998, Ladich and Yan 1998, Yan 1998, Yan and Curtsinger 2000, Yan et al. 2000, Scholik and Yan 2001). The ABR is a non-invasive far-field recording technique which acquires brainwaves generated from a series of neurogenerators upon the reception of acoustic stimuli from 1st-order receptors (e.g., sensory hair cells of the inner ear) along ascending auditory pathways to higher brain centers (e.g., the auditory cortex). The acoustically evoked brainwaves from bluegill sunfish (*Lepomis macrochirus*) (Rigedon and Yan 1997), goldfish (*Carassius auratus*), the oscar (*Astronotus ocellatus*) (Kenyon et al. 1998), croaking gourami (*Trichopsis vittata*), pygmy gourami (*Trichopsis pumila*), dwarf gourami (*Colisa lalia*), paradise fish (*Marcopodus opercularis*), blue gourami (*Trichogaster trichopterus*) (Ladich and Yan 1998), kissing gourami (*Helostoma temminckii*) (Yan 1998), toadfish (*Opsanus tau*) (Yan et al. 2000), mormyrid weakly electric fish (*Brienomyrus brachyistius*) (Yan and Curtsinger 2000), and fathead minnow (*Pimephales promelas*) (Scholik and Yan 2001) have been demonstrated to be species-specific (Yan et al. 1997). Each acoustically evoked brainwave indicates the summation of

neuronal activities of neurons along ascending auditory pathways, and the species specificity of the brainwave reflects the unique architecture of the neuronal network involved in the generation of acoustically evoked brainwaves. The 1st peak of the waveform is generated from the 3 hearing endorgans on each side of the inner ear and is also species-specific among many mammalian species examined (Hall 1992). It is known that taxonomical characters are variations of a homologous structure, and to be useful, they must show some variations in the taxon under study (Helfman et al. 1997). The species specificity of even the 1st peak of the acoustically evoked brainwave points to the possibility that the relative distribution of the otolithic organs of the fish may be species-specific and hence can be used as a valid character for use in systematic identification.

Otoliths of fish larvae are small and are relatively transparent under examination with a conventional dissecting microscope, which renders localization of these tiny otoliths difficult. Because of their calcareous nature and amino acid composition, they possess birefringence properties. With the aid of rotating polarization filters, transparent otoliths can be made to shine brightly against a darkened background when the phase retardation is set to $1/2\lambda$ (λ : wavelength of light used) (Slayter 1970, Rawlins 1992). Such a simple arrangement of polarization microscopy makes it easy to localize and to map the relative distribution of otoliths.

The purpose of this study was to test the hypothesis that the relative distribution of otoliths of larval fish can be used as a valid character for identification use. The application of double-polarization microscopy also helps to accelerate the mapping and measuring of the distribution of otoliths, hence shortening the time needed for species identification. Eight species of larval fish of a very divergent group of 4 families and 8 genera were compared to test the hypothesis.

MATERIALS AND METHODS

Fish larvae were collected from waters of Taiwan and Texas (see Table 1 for species, family, sampling sites, and sample size). Upon collection, the specimens were preserved in 70% ethyl alcohol. The trypsin enzyme method (Taylor 1967) was used to clear the specimens to facilitate the visualization of otoliths. Cleared specimens were examined by double-polarized microscopy as described below.

Double-polarization filter setup

A conventional dissecting microscope (Olympus SZX12) with a tungsten light source from underneath the stage was fitted with two 48-mm (diameter) rotary type polarized filters. The 1st polarizer was mounted directly underneath the glass stage (Fig. 1). The 2nd polarizer was mounted under the objective lens of the scope. The larval specimen was mounted onto a transparent Sylgard 184 silicone elastomer plate (Dow Corning, Midland, MI) with 00-size insect pins and immersed in 100% glycerin, i.e., the storage solution of cleared specimens. With the tungsten light on, the 2nd polarizer was rotated to align the incoming light which has passed through the 1st polarizer to yield a darkened background with brightened otoliths (Fig. 2a).

Comparison of the relative distribution of otoliths

The relative distances of 3 pairs of otoliths of each species were calculated by 3 indices: IUR = $(UU/IOD) \times 100\%$ (IUR: inter-utricular ratio; UU: distance between the 2 utricular otoliths, the lapillus; IOD: inter-orbital distance), ISR = $(SS/IOD) \times 100\%$ (ISR: inter-sacculus ratio; SS: distance between the 2 sacculus otoliths, the sagitta), ILR = $(LL/IOD) \times 100\%$ (ILR: inter-lagena ratio; LL: distance between the 2 lagena otoliths, the astericus) (see Fig. 2b for measurements). The IOD was used as a common denominator to standardize all indices measured, a common practice used for morphometric measurements in ichthyological studies (Helfman et al. 1997). All distance measurements were made with the aid of an ocular micrometer and were calibrated with a stage micrometer. Boxplot diagrams (Hubbs and Hubbs 1953) were used to illustrate the range, mean, and standard error of each index. The Wilcoxon two-sample test was used to compare the mean values of indices obtained between red drum, kingfish, and

banded drum. Due to limited sample sizes, no statistical test was applied to the remaining 5 fish species.

RESULTS

A total of 38 individuals representing 4 families, 8 genera, and 8 species was used in the study (Table 1). Most of the species have lapillus,




Fig. 1. Schematic diagram showing the optical path of double-polarized dissecting microscopy used to view larval otoliths. L: light source; R: reflecting mirror; P₁: 1st rotary polarizer; S: sample on top of a stage; P₂: 2nd rotary polarizer; O: objective lens; E: eyepiece. Arrows indicate the pathway of the light.

Table 1. Families, species, sampling sites, and numbers of individuals (sample size) of larvae used in the study

| Family | Species | Sampling site | Sample size |
|---------------|--------------------------------|------------------|-------------|
| Blennidae | <i>Istiblennius dussumieri</i> | Patozi, Taiwan | 3 |
| Leiognathidae | <i>Gazza minuta</i> | Tanshui, Taiwan | 2 |
| | <i>Leiognathus nuchalis</i> | Tanshui, Taiwan | 2 |
| Pomacentridae | <i>Abudefduf vaigiensis</i> | Patozi, Taiwan | 3 |
| Sciaenidae | <i>Cynoscion nebulosus</i> | Port Aransas, TX | 2 |
| | <i>Larimus fasciatus</i> | Port Aransas, TX | 6 |
| | <i>Menticirrhus americanus</i> | Port Aransas, TX | 5 |
| | <i>Sciaenops ocellatus</i> | Port Aransas, TX | 15 |

sagittae, and astericus otoliths, except for *Sciaenops ocellatus* and *Leiognathus nuchalis* in which the astericus (of laogenar chamber) could not be positively identified from the specimens examined, perhaps due to their small size.

Ranges of distances between the 3 pairs of otoliths (e.g., U-U, S-S, and L-L) are listed in Table 2. Relative distances between pairs of otoliths divided by the inter-orbital distance (IOD), i.e., the IUR and ISR indices, are shown as boxplot diagrams in figures 3 and 4, respectively. Due to limited sample

sizes on ILR data, no ILR boxplot diagram is provided. For the IUR index, the ratio in *S. ocellatus* has the highest mean value (109.8%) (Fig. 3, lower panel) while *Larimus fasciatus* has the lowest ratio (mean value: 53.61%) (Fig. 3, upper panel). When taking the 4 sciaenids as a group, the box diagrams (Fig. 3, lower panel) show that each species has its own unique mean IUR value. While the range of IUR values may overlap among species, e.g., between spotted seatrout and kingfish or kingfish and banded drum, there is no overlap between red drum and




Fig. 2. (a) Color photomicrograph showing an enzyme-cleared specimen (red drum) viewed under a double-polarization filter setup. Notice how the brightened birefringence properties of the otoliths against the darkened background make localization of the otoliths easy. U: utricular otolith; S: saccular otolith. Bar indicates 2 mm. (b) Schematic diagram showing measurements of distances made in this study. Distances were measured between eyes and 3 endorgans on each side from the ventral view of a larval fish skull. IOD: inter-orbital distance; U-U: distance between the 2 utricular otoliths; S-S: distance between the 2 saccular otoliths; L-L: distance between the 2 lagenar otoliths.




Fig. 3. Range (horizontal line), mean (vertical line) and 1 standard error (box) of inter-utricular distance/inter-orbital distance, i.e., IUR, of the species examined. Lower panel: red drum: *Sciaenops ocellatus*; spotted seatrout: *Cynoscion nebulosus*; kingfish: *Menticirrhus americanus*; banded drum: *Larimus fasciatus*. Upper panel: Toothpony *Gazza minuta*; Spotnape ponyfish *Leiognathus nuchalis*; blenny: *Istiblennius dussumieri*; damselfish: *Abudefduf vaigiensis*. The dotted line separates larvae collected from Taiwan (upper panel) and Texas (lower panel). Sample size refers to Table 1.

kingfish or red drum and banded drum. Results of the Wilcoxon two-sample test indicate that a significant difference is found between red drum and kingfish ($t_S = 7.52$, $P < 0.01$), red drum and banded drum ($t_S = 9.73$, $P < 0.01$), and between kingfish and banded drum ($t_S = 4.67$, $P < 0.01$). For species collected from Taiwan, *Gazza minuta* shows no overlap in IUR value with those of blenny (*Istiblennius dussumieri*) or damselfish (*Abudefduf vaigiensis*).

While *Gazza minuta* and *Leiognathus nuchalis* belong to the same family (Leiognathidae) and overlap in their IUR values, each species has its own unique mean IUR value (Fig. 3, upper panel).

For the ISR index, the ratio in *S. ocellatus* also ranks as the highest (mean value, 89.8%) (Fig. 4, lower panel), while *Istiblennius dussumieri* has the lowest value (mean, 43%) (Fig. 4, upper panel). Like that observed with IUR values of sciaenids, each

Table 2. Ranges of total length and distances between orbitals (IOD) and otoliths (utricle, U-U; saccule, S-S; and lagena, L-L)

| Family/Species | TL (mm) | IOD (mm) | U-U (mm) | S-S (mm) | L-L (mm) |
|--------------------------------|-------------|-----------|-----------|-----------|--------------|
| Blennidae | | | | | |
| <i>Istiblennius dussumieri</i> | 29.13-37.36 | 3.30-4.00 | 1.85-2.60 | 1.35-1.70 | 1.20-1.25 |
| Leiognathidae | | | | | |
| <i>Gazza minuta</i> | 15.89-17.07 | 1.75-1.80 | 1.50-1.60 | 0.90-1.00 | 0.75-0.80 |
| <i>Leiognathus nuchalis</i> | 10.42-12.00 | 1.32-1.45 | 0.75-1.28 | 0.80-1.10 | 0.65 |
| Pomacentridae | | | | | |
| <i>Abudefduf vaigiensis</i> | 20.30-29.47 | 3.00-4.40 | 1.50-2.60 | 1.20-1.90 | 1.10-1.50 |
| Sciaenidae | | | | | |
| <i>Cynoscion nebulosus</i> | 3.10-5.80 | 0.70-0.80 | 0.38-0.75 | 0.32-0.63 | 0.63 |
| <i>Larimus fasciatus</i> | 3.70-5.80 | 1.00-1.25 | 0.45-0.78 | 0.45-0.60 | 0.45-0.60 |
| <i>Menticirrhus americanus</i> | 3.10-6.00 | 0.95-1.50 | 0.60-0.75 | 0.40-0.96 | 0.64-0.80 |
| <i>Sciaenops ocellatus</i> | 6.40-9.00 | 0.50-0.75 | 0.55-0.90 | 0.50-0.65 | Not detected |




Fig. 4. Range (horizontal line), mean (vertical line), and 1 standard error (box) of inter-saccular distance/inter-orbital distance, i. e., ISR, of the species examined. Lower panel: red drum: *Sciaenops ocellatus*; spotted seatrout: *Cynoscion nebulosus*; kingfish: *Menticirrhus americanus*; banded drum: *Larimus fasciatus*. Upper panel: Toothpony: *Gazza minuta*; Spotnape ponyfish *Leiognathus nuchalis*; blenny: *Istiblennius dussumieri*; damselfish: *Abudefduf vaigiensis*. The dotted line separates larvae collected from Taiwan (upper panel) and Texas (lower panel). Sample size refers to Table 1.

sciaenid species has its own unique mean value of ISR (Fig. 4, lower panel). In terms of the range of ISR values, that of red drum does not overlap with that of banded drum, while it does overlap with the values of spotted seatrout and kingfish. Wilcoxon two-sample test results show significant differences in mean values between red drum and kingfish ($t_s = 6.34$, $P < 0.01$), red drum and banded drum ($t_s = 9.86$, $P < 0.01$), and kingfish and banded drum ($t_s = 4.32$, $P < 0.01$). The ISR values of *Gazza minuta* do not overlap with those of *Istiblennius dussumieri*, but overlap slightly with those of *Leiognathus nuchalis*. There is no overlap of ISR between *Istiblennius dussumieri* and *Leiognathus nuchalis* (Fig. 4, upper panel).

The IUR and ISR measurements of 4 sciaenids (red drum, spotted seatrout, kingfish, and banded drum) (lower panels of Figs. 3 and 4) and Wilcoxon two-sample test results clearly indicate that these 2 measurements are species-specific. Within the family Leiognathidae, the ISR value of *L. nuchalis* differs from that of *G. minuta*; while IUR values of these 2 species may overlap, the mean value of *G. minuta* is larger than that of *Leiognathus nuchalis*, another indication of species specificity of the spatial arrangement of otolithic organs. The IUR and ISR values of blenny (*I. dussumieri*) overlap with those of damselfish (*A. vaigiensis*), and using only these ratios as the sole character for identification may pose some difficulties. However, distinct differences in the external morphology between the 2 species easily minimize possible confusion.

DISCUSSION

Traditionally the identification of fish is based entirely or mainly on the morphological structures of adults (Hubbs and Lagler 1964). It was not until recently that early life history (ELH) stages have been used as main characters for systematics use (see review by Cohen 1984 and extensive work cited in Moser et al. 1984). The taxonomic characters used during embryonic stages include diameter of developing embryos, number and size of oil globules, distribution of oil globules, width of the perivitelline space, shape and size of the yolk, and degree of yolk segmentation (Moser and Ahlstrom 1970, Moser 1972, McGowan and Berry 1984). The main characters used for larval fish identification include: 1) meristic counts such as the number of fin rays, vertebrae, and myomeres (Gordon et al. 1984), and 2) morphometric measurements of larvae including the ratios of head length to body depth, head width

to head length, eye length to head length, and body depth to body length (Leis and Rennis 1983, Ahlstrom et al. 1984). Despite the usefulness of ELH in identifying larval species, the task itself is rather tedious and time consuming. Therefore, to identify a single key character that is species-specific would be of great value for quickly identifying targeted larval species for recruitment studies. The IUR and ISR indices developed in the present study provide an additional useful character for use in quick species identification. For instance, when both IUR and ISR ratios are taken into account, the 4 sciaenids can easily be separated by these values. This is even more valuable considering that all 4 sciaenids are found sympatrically and do not differ that much in their external morphology as well as body length. Cases like blenny and damselfish may overlap in their ratios; however, additional morphological characters such as a distinctive body shape and pigmentation patterns can aid in sorting out differences between species. A quick glance at the indices themselves also reveals the uniqueness of anatomical features of each species. For instance, both the IUR and ISR ratios of the red drum exceed 100% which clearly indicates that the cephalic region of the red drum is likely to have an anteriorly pointed triangular shape with inter-orbital distance narrower than the distance between inter-otolithic organs.

The advantage of using the indices developed herein is demonstrated in this study for quickly and accurately identifying larval fish. However, it is also imperative to point out that due to the nature of late formation of lagena otoliths (astericus), the ILR may not be as powerful as the values of IUR and ISR for identification use. The trypsin enzyme-clearing method was used in the present study to aid in easy visualization of otoliths, and it generally adds 1 additional day of effort for specimen preparation. However, for small-sized larvae (e.g., sciaenids: 3.1–9.0 mm), the location of otoliths can be identified easily even with just the use of double polarization filters without going through the enzyme-clearing process. The visualization of otoliths of larger size specimens, e.g., blennids (due to a thicker skull and muscle layers) especially can be greatly enhanced by clearing the samples.

In this study, only 2-dimensional distance values are measured and compared. The IUR, ISR, and ILR ratios used in the present study represent a simplified version of a Box-Truss measurement, which is a network of distance measures among homologous landmarks on a form, patterned as a series of contiguous quadrilaterals containing both internal diagonals (Brookstein et al. 1985). The full featured

Box-Truss method can be used to take into account distances among different otoliths and additional landmarks (e.g., eyes, tip of snout). However, in order to make the best use of the spatial distribution of otoliths as a powerful yet simple character for identification, a valid method of registering coordinates of 3-dimensional values of each otolith needs to be developed. This novel method also needs to incorporate a robust statistical method that can aid in discriminating the spatial distribution pattern of otoliths of each species. Future efforts should be directed to the development and refinement of registration of 3-dimensional spatial distribution of otoliths, so as to provide an even more powerful method of identifying larval fish. The limitation of sample size in 5 out of 8 species examined prevents further use of the *t*-test in comparing mean values between species. Nevertheless, significant statistical differences observed among 3 sciaenids clearly validate the merits of the methodology developed in the present study.

In conclusion, the data from this study support the hypothesis that the spatial distribution of otoliths is species-specific, and thus it can be used as a valid and practical character in larval fish identification. Furthermore, the use of readily available camera-type polarization filters is economically feasible and can provide a rather quick and convenient way of localizing otoliths in larval fish. The identification of larvae can become an easier task when used in conjunction with the indices developed in the study.

Acknowledgments: Dr. D. M. Checkley Jr. of the Scripps Oceanographic Institution provided technical guidance in the setup of double-polarization microscopy. Funding support to LSC was provided by the National Science Council of the R.O.C. (NSC89-2313-B-396-003). HYY was supported by grants from the United States National Institute of Mental Health (MH-58198), National Institute of Museum and Library Services (LL-90187), and National Organization for Hearing Research. Specimens of larval fish were provided by Dr. Y. Z. Wang of the Institute of Zoology, Academia Sinica, Dr. H. G. Moser of U.S. National Marine Fisheries Service, Southwest Fisheries Center, La Jolla, CA, and Dr. S. Holt and Mr. M. Li of the Univ. of Texas Marine Science Institute at Port Aransas, TX. Part of this work was presented in a poster format at the 2000 American Society for Ichthyologists and Herpetologists annual meeting in La Paz, Mexico. This work is dedicated to Prof. Clark Hubbs, Regents Emeritus Professor of the Univ. of Texas at Austin on his 80th birthday for his more than 60 yr of contributions

to ichthyology.

REFERENCES

- Ahlstrom EH, HG Moser, DM Cohen. 1984. Argentinoidae: development and relationships. *In* HG Moser, WJ Richards, DM Cohen, MP Fahay, AW Kendall Jr, SL Richardson, eds. *Ontogeny and systematics of fishes*. Amer. Soc. Ichthyol. Herpetol., Spec. Publ. 1, pp. 155-169.
- Bailey KM. 1982. The early life history of the Pacific hake *Merluccius products*. *Fish. B.-NOAA* **80**: 589-598.
- Bookstein F, B Chernoff, R Elder, J Humphries, G Smitj, R Strauss. 1985. *Morphometrics in evolutionary biology*. Philadelphia: Acad. Natl. Sci. Philadelphia, Spec. Publ. 15.
- Brothers EB. 1984. Otolith studies. *In* HG Moser, WJ Richards, DM Cohen, MP Fahay, AW Kendall Jr, SL Richardson, eds. *Ontogeny and systematics of fishes*. Amer. Soc. Ichthyol. Herpetol., Spec. Publ. 1, pp. 50-57.
- Brothers EB, CP Mathews, R Lasker. 1976. Daily growth increments in otolith from larval and adult fishes. *Fish. B.-NOAA* **74**: 1-8.
- Brothers EB, DM Williams, PF Sale. 1983. Length of larval life in 12 families of fishes at One Tree Lagoon, Great Barrier Reef, Australia. *Mar. Biol.* **76**: 319-324.
- Casteel RW. 1974. Identification of the species of Pacific salmon (Genus *Oncorhynchus*) native to North America based on otoliths. *Copeia* **1974**: 305-311.
- Chang CW, WN Tzeng, YC Lee. 2000. Recruitment and hatching dates of grey mullet (*Mugil cephalus* L.) juveniles in the Tanshui estuary of northwest Taiwan. *Zool. Stud.* **39**: 99-106.
- Chao LN. 1978. A basis for classifying Western Atlantic Sciaenidae (Teleostei: Perciforms). Washington, D.C. NOAA Tech Rept. NMFS Cir. 415.
- Chen KC, KT Shao, JS Yang. 1999. Using micropylar ultrastructure for species identification and phylogenetic inference among four species of Sparidae. *J. Fish Biol.* **55**: 288-230.
- Cheng PW, WN Tzeng. 1996. Timing of metamorphosis and estuarine arrival across the dispersal range of the Japanese eel *Anguilla japonica*. *Mar. Ecol. Prog. Ser.* **131**: 87-96.
- Claramunt RM, DH Wahl. 2000. The effects of abiotic and biotic factors in determining larval fish growth rates: a comparison across species and reservoirs. *Trans. Amer. Fish. Soc.* **129**: 835-851.
- Cohen DM. 1984. Ontogeny, systematics, and phylogeny. *In* HG Moser, WJ Richards, DM Cohen, MP Fahay, AW Kendall Jr, SL Richardson, eds. *Ontogeny and systematics of fishes*. Amer. Soc. Ichthyol. Herpetol., Spec. Publ. 1, pp. 7-11.
- Degens ET, WG Deuser, RL Haedrich. 1969. Molecular structure and composition of fish otoliths. *Mar. Biol.* **2**: 105-113.
- Fischer P. 1999. Otolith microstructure during the pelagic, settlement and benthic phases in burbot. *J. Fish Biol.* **54**: 1231-1243.
- Gallego A, MR Heath, DJ Basford, BR Mackenzie. 1999. Variability in growth rates of larval haddock in the northern North Sea. *Fish. Oceanogr.* **8**: 77-92.
- Gauldie RW, EJ Garynoth, J Illingworth. 1980. The relationship of the iron content of some fish otoliths to temperature.

- Comp. Biochem. Phys. **66**: 19-24.
- Gordon DJ, DF Markel, JE Olney. 1984. Ophidiiformes: development and relationships. In HG Moser, WJ Richards, DM Cohen, MP Fahay, AW Kendall Jr, SL Richardson, eds. Ontogeny and systematics of fishes. Amer. Soc. Ichthyol. Herpetol., Spec. Publ. 1, pp. 308-319.
- Hall JW. 1992. Handbook of auditory evoked responses. Boston: Allyn and Bacon.
- Hecht Y. 1978. A descriptive systematic study of the otoliths of the neopterygian marine fishes of South Africa. Part I. Introduction. Trans. Roy. Soc. S. Afr. **43**: 191.
- Helfman GS, BS Collette, DE Facey. 1997. The diversity of fishes. Malden, MA: Blackwell Science.
- Hubbs CL, C Hubbs. 1953. An improved graphical analysis and comparison of series of samples. Syst. Zool. **8**: 50-56.
- Hubbs CL, KF Lagler. 1964. Fishes of the Great Lakes region. Ann Arbor, MI: Univ. of Michigan Press.
- Huyghebaert B, D Nolf. 1979. An annotated bibliography of paleontological and systematic papers on fish-otoliths, published since 1968. Meded. Werkr. Tert. Kwart. Geol. **16**: 139-170.
- Kendall AW Jr, D Gordon. 1981. Growth rate of Atlantic mackerel (*Scomber scombrus*) larvae in the middle Atlantic Bight. Rapp. P. V. Reun. Cons. Int. explor. Mer. **178**: 337-341.
- Kenyon TN, F Ladich, HY Yan. 1998. A comparative study of hearing ability in fishes: the auditory brainstem response approach. J. Comp. Phys. A **182**: 307-318.
- Kokita T, M Omori. 1999. Long distance dispersal of larval and juvenile rockfish, *Sebastes thompsoni*, with drifting seaweed in the Tohoku area, northwest Pacific, estimated by analysis of otolith microstructure. B. Mar. Sci. **65**: 105-118.
- Ladich F, HY Yan. 1998. Correlation between auditory sensitivity and vocalization in anabantoid fishes. J. Comp. Phys. A **182**: 737-746.
- Laroche JL, S Richardson, AA Rosenberg. 1982. Age and growth of a pleuronectid, *Parophrys vetulus*, during the pelagic larval period in Oregon coastal waters. Fish. B.-NOAA **75**: 547-554.
- Leis JM, DS Rennis. 1983. The larvae of Indo-Pacific coral reef fishes. Sydney, Australia: New South Wales Univ. Press.
- Lou DC, NA Moltschanidkyj. 1992. Daily otolith increments in juvenile tropical parrotfishes and surgeonfishes. Aust. J. Mar. Freshwater Res. **43**: 974-981.
- Lough RG, M Pennington, GR Bolz, AA Rosengerg. 1982. Age and growth of larval Atlantic herring, *Clupea harengus* L., in the Gulf of Maine-Georgia Bank region based on otolith growth increments. Fish. B.-NOAA US **80**: 187-199.
- Lowenstan HA. 1980. Bioinorganic constituents of hard parts. In PE Hare, ed. Biogeochemistry of amino acids. New York: Wiley, pp. 3-16.
- Lowenstan HA. 1981. Minerals formed by organisms. Science **211**: 1126-1131.
- Lowenstan HA, JE Fitch. 1978. Morphology, mineral diversity and ontogenetic mineralogic changes in extant elasmobranch otoconia. Geol. Soc. Am. Abstr. **10**: 114.
- Lowenstan HA, JE Fitch. 1981. Vaterite formation in otoliths of higher teleost fishes. Abstract, Pacific Division, AAAS, Annual Meeting, 14-19 June 1981. Univ. Oregon, Eugene.
- McGowan MF, FH Berry. 1984. Clupeiformes: development and relationships. In HG Moser, WJ Richards, DM Cohen, MP Fahay, AW Kendall Jr, SL Richardson, eds. Ontogeny and systematics of fishes. Amer. Soc. Ichthyol. Herpetol., Spec. Publ. 1, pp. 108-126.
- Methot RD, D Kramer. 1979. Growth of northern anchovy, *Engraulis mordax*, larvae in the sea. Fish. B.-NOAA **77**: 1124-1127.
- Moser HG. 1972. Development and distribution of the rockfish, *Sebastes macdonaldi* (Eigenmann and Beeson, 1893), family Scorpaenidae off southern California and Baja California. Fish. B.-NOAA **70**: 941-958.
- Moser HG, EH Ahlstrom. 1970. Development of lanternfishes (Family Myctophidae) in the California Current. Part I. Species with narrow-eyed larvae. B. LA. Cty. Mus. Nat. Hist. Sci. **7**, 145 pp.
- Moser HG, EH Ahlstrom. 1974. Role of larval stages in systematic investigations of marine teleosts: the Myctophidae, a case study. Fish. B.-NOAA US **72**: 391-413.
- Moser HG, WJ Richards, DM Cohen, MP Fahay, AW Kendall Jr, SL Richardson, eds. 1984. Ontogeny and systematics of fishes. Amer. Soc. Ichthyol. Herpetol., Spec. Publ. 1.
- Panella AL. 1971. Fish otoliths: daily growth layers and periodical patterns. Science **173**: 1124-1127.
- Panella AL. 1980. Growth patterns in fish sagittae. In Rhoads DC, RA Lutz, eds. Skeletal growth of aquatic organisms; biological records of environmental change. New York: Plenum Press, pp. 519-560.
- Papadopoulou C, GD Kanis, E Moraitopoulou-Kassimati. 1978. Zinc content in otoliths of mackerel from the Aegean. Mar. Pollut. Bull. **9**: 106-108.
- Papadopoulou C, GD Kanis, E Moraitopoulou-Kassimati. 1980. Trace element content in fish otoliths in relation to age and size. Mar. Pollut. Bull. **11**: 68-72.
- Paperno R, TE Targett, PA Grecay. 2000. Spatial and temporal variation in recent growth, overall growth, and mortality of juvenile weakfish (*Cynoscion regalis*) in Delaware Bay. Estuaries **23**: 10-20.
- Rawlings DJ. 1992. Light microscopy. Oxford: BIOS Sci. Publishers.
- Rigdon SM, HY Yan. 1997. Comparisons of auditory sensitivities and sensory hair cell densities between adult and juvenile bluegill sunfish. New Orleans: Soc. Neurosci. Ann. Meeting, Abstract 77.11.
- Rooker JR, SA Holt, JG Holt, LA Fuiman. 1999. Spatial and temporal variability in growth, mortality, and recruitment potential of postsettlement red drum, *Sciaenops ocellatus*, in a subtropical estuary. Fish. B.-NOAA **97**: 581-590.
- Scholik AR, HY Yan. 2001. Effects of underwater noise on auditory sensitivity of a cyprinid fish. Hearing Res. **152**: 17-24.
- Slayer EM. 1970. Optical methods in biology. New York: Wiley-Interscience.
- Struhsaker P, JH Uchiyama. 1976. Age and growth of the nehu *Stolephorus purpureus* (Pisces: Engraulidae) from the Hawaiian Islands as indicated by daily growth increments of sagittae. Fish. B.-NOAA **74**: 9-17.
- Taylor WR. 1967. An enzyme method of clearing and staining small vertebrates. Proc. US Nat. Mus. **122**: 1-17.
- Thorrold SR, CM Jones, PL Swart, TE Targett. 1998. Accurate classification of juvenile weakfish *Cynoscion regalis* to estuarine nursery area based on chemical signatures in otoliths. Mar. Ecol. Prog. Ser. **173**: 253-265.
- Townsend DW, JJ Graham. 1981. Growth and age structure of larval Atlantic herring, *Clupea harengus*, in the Sheepscot River estuary, Maine, as determined by daily growth increments in otoliths. Fish. B.-NOAA. **79**: 123-

- 130.
- Tzeng WN. 1995a. Migratory history recorded in otoliths of the Japanese eel, *Anguilla japonica*, elvers as revealed from SEM and WDS analyses. Zool. Stud. **34** (Supplement 1): 234-236.
- Tzeng WN. 1995b. Temperature effects on the incorporation of strontium in otolith of Japanese eel, *Anguilla japonica*. J. Fish Biol. **45**: 1055-1066.
- Tzeng WN. 1996. Effect of salinity and ontogenetic movement on Sr/Ca ratio in otoliths of the Japanese eel, *Anguilla japonica* Temminck & Schlegel. J. Exp. Mar. Biol. Ecol. **199**: 111-122.
- Tzeng WN, YC Tsai. 1994. Changes in otolith microchemistry of the Japanese eel, *Anguilla japonica*, during its migration from the ocean to the rivers of Taiwan. J. Fish Biol. **45**: 671-684.
- Tzeng WN, HF Wu, H Wickstrm. 1994. Scanning electron microscopic analysis of annulus microstructure in otolith of European eel, *Anguilla anguilla*. J. Fish Biol. **45**: 479-492.
- Wang CH, WN Tzeng. 1999. Differences in growth rates among cohorts of *Encrasicholina punctifer* and *Engraulis japonicus* larvae in the coastal waters off Tanshui River Estuary, Taiwan, as indicated by otolith microstructure analysis. J. Fish Biol. **54**: 1002-1016.
- Wang CH, WN Tzeng. 2000. The timing of metamorphosis and growth rates of American and European eel leptocephali: a mechanism of larval segregative migration. Fish. Res. **46**: 191-205.
- Wilson DT, MI McCormick. 1997. Spatial and temporal validation of settlement-marks in the otoliths of tropical reef fishes. Mar. Ecol. Prog. Ser. **153**: 259-271.
- Wilson DT, MI McCormick. 1999. Microstructure of settlement-marks in the otoliths of tropical reef fishes. Mar. Biol. **134**: 29-41.
- Yan HY. 1998. Auditory role of the suprabranchial chamber in gourami fish. J. Comp. Phys. A **183**: 325-333.
- Yan HY, WS Curtsinger. 2000. The otic gasbladder as an ancillary auditory structure in a mormyrid fish. J. Comp. Phys. A **186**: 595-602.
- Yan HY, ML Fine, NS Horn, WE Colón. 2000. Variability in the role of the gasbladder in fish audition. J. Comp. Phys. A **186**: 435-445.
- Yan HY, SM Rigdon, A Dutschke. 1997. Auditory brainstem response brainwaves of fish are species specific. New Orleans: Soc. Neurosci. Ann. Meeting, Abstract 77.10.
- Zhang Z, RJ Beamish. 2000. Use of otolith microstructure to study history of juvenile chinook salmon in the Strait of Georgia in 1995 and 1996. Fish. Res. **46**: 239-250.

以耳石的相對位置當作仔魚鑑定的形質

陳麗淑¹ 嚴宏洋²

最近發展的聽覺腦幹反應(ABR)技術指出, 魚類聽覺引發的腦波具有種的特異性。這些腦波反映聽覺核和沿著往上的聽覺神經通路的空間分布。從腦波的種特異性可以引申出耳石的分布應該也是具有種特異性的假說, 因為耳石由面所在的感覺毛激發的反應引發 ABR 第一個波。本研究從臺灣和美國德州採用八種仔魚(分屬於四科八屬), 以胰蛋白酶將魚體處理成透明, 再用解剖顯微鏡裝置偏光鏡觀察。三個參數(囊狀石間距 inter-utricle ratio, IUR; 矢狀(扁平)石間距 inter-sacculus ratio, ISR; 星狀石間距 inter-lagena ratio, ILR)用來度量耳石間距和眼距的比例。IUR 和 ISR 可以成功地把種分開, 但 ILR 因為星石發育較晚就顯不出作用。本研究結果支持原先所提的假說: 仔魚耳石的空間分布可以當成一個有效的形質來分離不同的種。

關鍵詞: 機械式感覺生理學, 內耳, 耳石, 聽覺路徑, 分類。

¹ 國立海洋科技博物館籌備處

² School of Biological Sciences, University of Kentucky, Lexington, KY 40506-0225, USA