

Erratum

Hematological Profiles of the Formosan Black Bear (*Ursus thibetanus-formosanus*)

Geng-Ruei Chang^{1,*}, Frank Chiahung Mao¹, Chieh-Chung Yang², and Fang-Tse Chan²

¹Department of Veterinary Medicine, National Chung Hsing University, 250 Kuo-Kuang Road, Taichung, Taiwan 402, R.O.C.

²Endemic Species Research Institute, Council of Agriculture, 1 Ming-Sheng E. Road, Chichi, Nantou, Taiwan 552, R.O.C.

[Article in *Zoological Studies* 45(1): 93-97(2006)]

In page 94, a paragraph of the RESULT has been misprinted. They should be shown as below.

RBC, Hb, Hct, and MCV showed significant differences ($p < 0.05$) between seasons, while differences between sexes were not significant (Table 1). Results indicated a significant increase in autumn-winter compared with values in spring-summer. Female WBC and lymphocyte counts were significantly higher ($p < 0.05$) in spring-summer than in autumn-winter. Monocyte counts illustrated the opposite response (Table 2). Differences between sexes were only found to be significant for MCH values ($p < 0.05$).

In Table 1. Comparison of hematological values (mean \pm SE, $n = 21$, 10 each for females and males) between sexes and season for adult Formosan black bears has been misprinted. They should be shown as below.

Table 1. Comparison of hematological values (mean \pm SE, $n = 21$, 10 each for females and males) between sexes and season for adult Formosan black bears

Blood parameter	Sex	Season	
		Spring-summer	Autumn-winter
RBC ^a ($10^6/\mu\text{l}$)	F	5.96 \pm 0.16	6.80 \pm 0.18
	M	6.35 \pm 0.18	7.15 \pm 0.50
Hb ^a (g/dl)	F	13.93 \pm 0.33	15.58 \pm 0.43
	M	14.90 \pm 0.09	17.96 \pm 1.01
Hct ^a (%)	F	38.80 \pm 0.98	44.60 \pm 1.68
	M	41.60 \pm 1.55	49.00 \pm 2.17
MCV ^a (fl)	F	61.90 \pm 1.06	66.00 \pm 1.25
	M	61.70 \pm 3.11	69.60 \pm 1.76
MCH ^b (pg)	F	22.00 \pm 0.49	23.20 \pm 0.09
	M	23.60 \pm 0.31	24.30 \pm 0.07
MCHC (g/dl)	F	35.31 \pm 0.91	35.60 \pm 0.35
	M	34.60 \pm 0.31	35.80 \pm 0.10
PLT ($10^3/\mu\text{l}$)	F	196.00 \pm 42.30	229.00 \pm 47.70
	M	311.00 \pm 22.40	258.00 \pm 65.20

^aBlood values significantly differed ($p < 0.05$) between seasons.

^bBlood values significantly differed ($p < 0.05$) between sexes. RBC, total erythrocyte count; Hb, hemoglobin; Hct, hematocrit; MCV, mean corpuscular volume; MCH, mean corpuscular hemoglobin; MCHC, mean corpuscular hemoglobin concentration; PLT, total platelet count.

In Table 2. Comparison of leucocyte counts (mean \pm SE, $n = 22$, 7 each for females and males) and differential counts of leucocytes in percentage (mean \pm SE, $n = 16$, 7 each for females and males) between sexes for adult Formosan black bears has been misprinted. They should be shown as below.

Table 2. Comparison of leucocyte counts (mean \pm SE, $n = 22$, 7 each for females and males) and differential counts of leucocytes in percentage (mean \pm SE, $n = 16$, 7 each for females and males) between sexes for adult Formosan black bears

Blood parameter	Sex	Season	
		Spring-summer	Autumn-winter
Total leucocyte counts ($10^3/\mu\text{l}$)	F	10.20 \pm 0.51	7.81 \pm 0.69*
	M	ND	8.33 \pm 1.19
Band cells (%)	F	1.00 \pm 0.55	1.20 \pm 0.39
	M	ND	5.60 \pm 2.50
Segmented cells (%)	F	62.25 \pm 5.82	62.20 \pm 4.12
	M	ND	55.00 \pm 3.78
Eosinophil cells (%)	F	0	5.30 \pm 1.60
	M	ND	7.60 \pm 2.23
Basophile cells (%)	F	0	0
	M	ND	0.20 \pm 0.20
Lymphocytes (%)	F	42.80 \pm 8.87	20.30 \pm 3.50*
	M	ND	22.20 \pm 5.31
Monocytes (%)	F	3.20 \pm 0.86	10.80 \pm 2.67*
	M	ND	9.40 \pm 3.36

*Female blood values significantly differed ($p < 0.05$) among seasons. ND, No samples were determined.