

Allozyme Variation in the Pacific Oyster *Crassostrea gigas* along the Coast of Taiwan

Li-Lian Liu^{1,*}, Keryea Soong¹ and Chang-Po Chen²

¹Institute of Marine Biology, National Sun Yat-sen University, Kaohsiung, Taiwan 804, R.O.C.

²Institute of Zoology, Academia Sinica, Nankang, Taipei, Taiwan 115, R.O.C.

(Accepted March 28, 1995)

Li-Lian Liu, Keryea Soong and Chang-Po Chen (1995) Allozyme variation in the Pacific oyster *Crassostrea gigas* along the coast of Taiwan. *Zoological Studies* 34(3): 177-182. Nine populations of the oyster *Crassostrea gigas*, were assayed for allozyme variation by starch gel electrophoresis. Three of the seven enzyme loci investigated are polymorphic: *Lap*, *Mpi*, and *Pgi*. Significant heterozygote deficiency was found in all seven Taiwan populations but not in the two populations of the Penghu Islands. Low levels of genetic variability ($p = 0.14 - 0.42$; $H = 0.044 - 0.084$) and little genetic differentiation (Nei's $D < 0.01$) were found among populations. Individuals with *Lap* heterozygotes were smaller than those with homozygotes in one population ($p = 0.03$) but this difference was statistically insignificant when adjusted significance level for multiple tests was used. Our results suggest that environmental factors may be important in determining regional differences in oyster size and shape.

Key words: Genetic variability, Heterozygosity.

The Pacific oyster *Crassostrea gigas* (Thunberg) ranges from the coast of Japan and the Korean peninsula to China and Southeast Asia (Ahmed 1975). It has also been introduced into Australia, France, New Zealand, and North America for commercial farming (Chew 1990). *C. gigas* is one of the most important cultured shellfish along the west coast of Taiwan and the adjacent Penghu Islands (Lin and Liang 1982).

In recent years, it has been reported that allozyme heterozygosity is positively correlated with growth rate in some marine bivalves, e.g., Pacific oyster *C. gigas* (Fujio 1982), American oyster *C. virginica* (Singh and Zouros 1978, Zouros et al. 1980), and blue mussel *Mytilus edulis* (Koehn and Gaffney 1984). There is little information about genetic variation and growth rate in *C. gigas* from Taiwan but regional differences in size and shape among aquaculture sites were reported by Soong et al. (1992). This study was undertaken to determine: (1) whether genetic variation exists among oysters from various locations; and (2) whether there is

any correlation between heterozygosity and size (expressed as shell length and wet meat weight) in *C. gigas* at collection sites in Taiwan.

MATERIALS AND METHODS

Oysters were collected along the west coast of Taiwan and the Penghu Islands from September 1990 to January 1991, from seven aquaculture areas at Shunsun, Lukang, Dongshi, Budai, Chigu, Dongkang, and Tsaiyuan, and two wild sites at Yunan and Waton (Fig. 1). Three to five long lines with adult-sized oysters attached to artificial substrata were sampled at each aquaculture site. Oysters at wild sites were collected from natural substrata. Live oysters were stored at -70°C after transport back to the laboratory. After shell length and wet meat weight were measured, mantle tissues were homogenized in a Tekmar tissumizer with an equal volume of 10 mM Tris-HCl buffer (pH 7.0) containing 1% Triton X-100. Homogenates

*To whom all correspondence and reprint requests should be addressed.




Fig. 1. *Crassostrea gigas* sample locations in Taiwan and the Penghu Islands.

were centrifuged at 5,000 g for 15 min and the supernatants were again stored at -70°C until electrophoretic analysis.

Seven loci were assayed: adenylate kinase (*Adk*, EC 2.7.4.3), arginine kinase (*Ark*, EC 2.7.3.3), isocitrate dehydrogenase (*Idh*, EC 1.1.1.42), leucine aminopeptidase (*Lap*, EC 3.4.11 or 13), malate dehydrogenase (*Mdh*, EC 1.1.1.37), mannose-6-phosphate isomerase (*Mpi*, EC 5.3.1.8), and phosphoglucoisomerase (*Pgi*, EC 5.3.1.9). Horizontal starch-gel electrophoresis with Tris-citrate pH 7.0 and 8.0 buffer systems were used. Enzyme-staining methods followed Murphy et al. (1990).

Alleles at each locus were scored by designating the most common allele as 100. All other alleles were numbered according to their relative anodal distance from the most common allele. Chi-square goodness-of-fit tests were computed at each locus to determine if there were significant deviations from Hardy-Weinberg equilibrium be-

tween observed and expected heterozygous genotype frequencies. The mean observed and expected heterozygosity in each population was also calculated. For each locus, the nine populations were tested for homogeneity using the G-contingency test (Sokal and Rohlf 1981) with pooled homozygotes and pooled heterozygotes. Nei's genetic distance coefficients (D) were also calculated. The Mann-Whitney U test was used to determine whether there was any difference between genotypes (homozygotes vs. heterozygotes) and size (expressed as shell length or wet meat weight).

RESULTS

When the commonest allele ≤ 0.95 was used as the criterion for polymorphism, four of the seven enzyme loci investigated were monomorphic: *Adk*, *Ark*, *Idh*, and *Mdh*. Allelic frequencies of polymorphic loci in the surveyed populations are shown in Table 1. Observed heterozygosity in the polymorphic *Lap*, *Mpi*, and *Pgi* loci among populations were 0.125 – 0.292, 0.028 – 0.227, and 0.050 – 0.181, respectively. Significant heterozygote deficiency was found at two to three polymorphic loci in all seven populations from Taiwan but not in the two populations from Penghu (Tsaiyuan and Waton). Mean heterozygosity of all loci in each population ranged from 0.044 to 0.084 and no significant difference was found between cultured and wild populations (Table 2).

The nine populations were homogeneous at *Lap* and *Pgi* and heterogeneous at *Mpi* ($G_{(8)} = 20.2$, $p < 0.01$). Nei's genetic distances among populations ranged from 0 to 0.004. In other words, very little genetic differentiation occurs at these seven loci among oyster populations in Taiwan and Penghu.

Due to low frequencies of heterozygotes found at the *Mpi* and *Pgi* loci, the relationship between heterozygosity and size was only assessed at the *Lap* locus. We found that, in the Budai population, individuals of *Lap* heterozygotes were significantly smaller than those of homozygotes ($p = 0.03$) (Table 3). However, by using adjusted significance level for multiple tests, this difference was statistically insignificant.

DISCUSSION

Populations of *C. gigas* in Taiwan and Penghu have lower mean heterozygosity (0.044 – 0.084)

Table 1. Allelic frequencies for three polymorphic loci in nine populations of the oyster *Crassostrea gigas*

Locus	Allele (RM)	Population								
		Cultured							Wild	
		Shunsun	Lukang	Dongshi	Budai	Chigu	Dongkang	Tsaiyuan	Yunan	Watou
<i>Lap</i>	(N)	72	72	68	72	117	79	66	44	59
	125	.000	.000	.000	.000	.013	.000	.000	.034	.000
	120	.014	.000	.000	.000	.000	.000	.000	.000	.000
	113	.097	.042	.015	.028	.056	.051	.015	.034	.068
	100	.819	.868	.919	.924	.825	.823	.886	.795	.864
	90	.007	.000	.000	.000	.021	.006	.000	.000	.000
	88	.049	.090	.066	.049	.077	.114	.091	.136	.034
	78	.014	.000	.000	.000	.009	.006	.008	.000	.034
	<i>Ho</i>	.292**	.181**	.132	.125	.282**	.203**	.197	.227**	.254
	<i>He</i>	.318	.238	.152	.145	.311	.309	.207	.350	.248
<i>Mpi</i>	(N)	72	72	69	72	118	79	66	48	60
	113	.000	.021	.029	.014	.000	.000	.000	.010	.008
	100	.986	.917	.949	.944	.936	.937	.886	.906	.983
	90	.000	.000	.000	.000	.000	.000	.000	.010	.000
	75	.014	.056	.022	.042	.064	.063	.114	.063	.008
	67	.000	.007	.000	.000	.000	.000	.000	.010	.000
	<i>Ho</i>	.028	.125**	.072**	.083*	.093**	.101	.227	.104**	.033
	<i>He</i>	.028	.157	.098	.107	.120	.119	.203	.176	.033
<i>Pgi</i>	(N)	72	72	69	72	118	76	66	44	60
	143	.090	.021	.022	.042	.064	.020	.053	.000	.000
	100	.889	.896	.957	.938	.886	.901	.924	.875	.975
	56	.021	.083	.022	.021	.051	.072	.023	.125	.017
	47	.000	.000	.000	.000	.000	.007	.000	.000	.008
	<i>Ho</i>	.083**	.181	.072**	.097*	.161**	.171**	.152	.068**	.050
	<i>He</i>	.203	.191	.085	.120	.210	.183	.144	.221	.049

(RM: relative allelic mobility; N: sample size; *Ho*, *He*: observed and expected heterozygosity from Hardy-Weinberg equilibrium; deviations from expected frequencies are tested by the X^2 -test. **: $p < 0.01$, *: $p < 0.05$).

Table 2. Summary of genetic variation in *Crassostrea gigas* in Taiwan

Population	Mean number alleles per locus \pm S.E.	Percent polymorphic loci (95% criterion)	Mean heterozygosity \pm S.E.	
			observed	expected
Cultured				
Shunsun	2.1 \pm 0.7	28.6	0.058 \pm 0.041	0.078 \pm 0.049
Lukang	2.1 \pm 0.5	42.9	0.071 \pm 0.033	0.086 \pm 0.040
Dongshi	2.1 \pm 0.3	28.6	0.044 \pm 0.019	0.052 \pm 0.023
Budai	2.1 \pm 0.4	42.9	0.050 \pm 0.020	0.059 \pm 0.024
Chigu	2.4 \pm 0.6	42.9	0.079 \pm 0.041	0.094 \pm 0.047
Dongkang	2.3 \pm 0.6	42.9	0.070 \pm 0.033	0.093 \pm 0.045
Tsaiyuai	2.0 \pm 0.4	42.9	0.084 \pm 0.039	0.081 \pm 0.037
Wild				
Yunan	2.1 \pm 0.6	42.9	0.057 \pm 0.032	0.107 \pm 0.054
Watou	2.0 \pm 0.5	14.3	0.048 \pm 0.035	0.047 \pm 0.034

Table 3. Average shell length and meat weight of heterozygotes and homozygotes at *Lap* locus in *Crassostrea gigas*

Population	N	Length (mm)		Weight (g)		
		Heterozygotes	N	Homozygotes	Heterozygotes	Homozygotes
Shunsun	21	38.3 ± 9.6	51	36.2 ± 7.4	6.05 ± 3.30	5.04 ± 2.57
Lukang	13	34.6 ± 7.9	58	37.6 ± 9.8	5.14 ± 3.57	6.00 ± 3.50
Dongshi	9	45.4 ± 17.7	57	43.6 ± 10.7	6.88 ± 4.11	7.74 ± 5.59
Budai	9	24.4 ± 14.9	63	36.2 ± 14.4*	1.88 ± 2.61	5.33 ± 4.52*
Chigu	33	43.5 ± 9.3	84	43.9 ± 8.3	5.02 ± 1.78	5.09 ± 1.83
Dongkang	16	34.4 ± 8.0	63	33.9 ± 9.1	5.70 ± 2.96	4.67 ± 3.25
Tsaiyuan	13	59.2 ± 10.6	53	59.4 ± 12.5	12.68 ± 5.70	11.34 ± 6.12

Data are presented as mean ± std; N: sample size; significance of the difference is given (*: $p < 0.05$).

than conspecific populations in other areas and other *Crassostrea* species. For instance, the mean heterozygosities were 0.186 (Fujio et al. 1983) and 0.234 (Ozaki and Fujio 1985) in *C. gigas* and *C. angulata*, respectively. In locus by locus comparisons, the observed heterozygosities of most loci in the surveyed populations were generally lower than other *C. gigas* populations, although the comparison was only based on six loci. For example, observed heterozygosities at *Lap* and *Pgi* were 0.125 – 0.292 and 0.050 – 0.181, respectively, in the present study, but 0.256–0.571 and 0.217–0.474 in the study of Buroker et al. (1979) in Japan. Many factors might result in the phenomenon of low heterozygosity in the populations in Taiwan and Penghu including the Wahlund effect, null alleles, inbreeding, small effective population size originally, bottleneck effect at Pleistocene due to sea level decrease (Emery et al. 1971), artificial selection, and others. However, it is not possible to elucidate the real cause at present.

The small genetic distances (0 to 0.004) among the nine populations of *C. gigas* indicate that little genetic differentiation occurs over a horizontal distance of up to 300 km. In Taiwan, oysters spawn throughout the year and the planktonic larval period ranges from one to two weeks (Lin and Liang 1982). Dongshi and Budai are the spat collection areas which supply spats for culturing in Taiwan and Penghu. The existence of planktonic larval stages and the procurement of spats from a single location has produced extensive gene flow which may have prevented genetic differentiation of these oyster populations. In addition, the practice by farmers of moving adult oysters between sites to increase their weight before marketing (personal observation) may also prevent genetic differentiation among populations.

Significant heterozygote deficiency was found in all populations from Taiwan but not in populations from Penghu. In the Penghu Islands, both cultured population (Tsaiyuan) and wild population (Waton) exist. The spats for culturing at Tsaiyuan are imported from Taiwan. Although extensive gene flow might diminish population differences between Taiwan and Penghu, heterozygote deficiency only occurs in Taiwan. The presence of null alleles, the Wahlund effect, and inbreeding, among other factors, are important in generating the phenomenon of heterozygote deficiency (Singh and Green 1984). If the presence of null alleles is an important factor, heterozygote deficiency should be observed in both Taiwan and Penghu. Inbreeding can not explain the phenomenon either, as cultured oysters are all from the same geographic origin. In this case, differential mortality due to local environmental factors might play a role in causing the genetic differences between adult oysters of Taiwan and Penghu.

Only the Budai population exhibited a size difference between oysters carrying heterozygous *Lap* genotypes (smaller) and those carrying homozygous *Lap* genotypes (larger). But this difference was not significant when adjusted significance level for multiple tests was used (Hochberg 1988). The relationship between heterozygosity and growth has been studied for various marine bivalves (Zouros and Pogson 1994). In the five-month-old blue mussel *Mytilus edulis*, a significant positive correlation was recorded between observed heterozygosity (total number of heterozygotes divided by sample size) and growth rate (samples separated into different size categories) at the octopine dehydrogenase locus (Gosling 1989). In the oyster *Ostrea edulis* (Alvarez et al. 1989), the relationship between heterozygosity and growth changed through time.

Positive correlations were found at isocitrate dehydrogenase-2 and malate dehydrogenase-1 loci in the eighteen-month-old oyster and at the isocitrate dehydrogenase-2 and esterase-3 loci in the thirty-month-old oyster. In the scallop *Placopecten magellanicus* (Volckaert and Zouros 1989), no significant difference was found between average shell height of heterozygotes and homozygotes at the 6-phosphogluconate dehydrogenase and glucose-phosphate isomerase loci. The relationship between heterozygosity and growth clearly depends on the enzyme locus studied, as well as the age of the organisms.

A study of American oyster *C. virginica* indicated that allozyme variation and shell morphology (i.e., shell shape and size) are not correlated (Groue and Lester 1982). Others have suggested that site-specific growth rates in *C. gigas* (Brown 1988) and in mussel *M. edulis* (Incze et al. 1980) are mostly influenced by environmental factors such as food supply, temperature, salinity and physiological conditions. Although age differences may also affect oyster size, the oysters at Tsaiyuan (Penghu) grow faster and larger than those cultured in Taiwan (Soong et al. 1992). With little genetic differentiation found among these oyster populations, regional differences in size and shape probably result from environmental factors.

Acknowledgements: We thank Messrs. Jau-Luren Chen and Dow-Yuan Soong, Misses Hwey-Lin Ho, Wei-Ming Liu, and Zon-Zon Tsai for assistance in field and laboratory work. We also thank Messrs. Wan-Sen Tsai and Chwen-Herng Wu for help collecting oysters in Penghu and Taishi. Three anonymous reviewers provided valuable advice to earlier versions of the manuscript. This study was sponsored by the Council of Agriculture of the Republic of China.

REFERENCES

- Ahmed M. 1975. Speciation in living oysters. *Advance in Mar. Biol.* **13**: 357-397.
- Alvarez G, C Zapata, R Amaro, A Guerra. 1989. Multilocus heterozygosity at protein loci and fitness in the European oyster, *Ostrea edulis* L. *Heredity* **63**: 359-372.
- Brown JR. 1988. Multivariate analysis of the role of environmental factors in seasonal and site-related growth variation in the Pacific oyster *Crassostrea gigas*. *Mar. Ecol. Prog. Ser.* **45**: 225-236.
- Buroker NE, WK Hershberger, KK Chew. 1979. Population genetics of the Family Ostreidae. I. Intraspecific studies of *Crassostrea gigas* and *Saccostrea commercialis*. *Mar. Biol.* **54**: 157-169.
- Chew KK. 1990. Global bivalve shellfish introductions. *World Aquacul.* **21**: 9-22.
- Emery KO, H Niino, B Sullivan. 1971. Post-Pleistocene levels of the East China Sea. Woods Hole Oceanographic Institute, Woods Hole, Massachusetts, Contribution no. **2441**: 381-390.
- Fujio Y. 1982. A correlation of heterozygosity with growth rate in the Pacific oyster, *Crassostrea gigas*. *Tohoku J. Agricul. Res.* **33**: 66-75.
- Fujio Y, R Yamanaka, PJ Smith. 1983. Genetic variation in marine molluscs. *Bull. Jpn. Soc. Sci. Fisher.* **49**: 1809-1817.
- Gosling EM. 1989. Genetic heterozygosity and growth rate in a cohort of *Mytilus edulis* from the Irish coast. *Mar. Biol.* **100**: 211-215.
- Groue KJ, LJ Lester. 1982. A morphological and genetic analysis of geographic variation among oysters in the Gulf of Mexico. *Veliger* **24**: 331-335.
- Hochberg Y. 1988. A sharper Bonferroni procedure for multiple tests of significance. *Biometrika* **75**: 800-802.
- Incze LS, RA Lutz, L Watling. 1980. Relationships between effects of environmental temperature and seston on growth and mortality of *Mytilus edulis* in a temperate Northern estuary. *Mar. Biol.* **57**: 147-156.
- Koehn RK, PM Gaffney. 1984. Genetic heterozygosity and growth rate in *Mytilus edulis*. *Mar. Biol.* **82**: 1-7.
- Lin YS, MH Liang. 1982. Growth and setting of cultured oyster (*Crassostrea gigas* Thunberg) in Budai Bay. *Bull. Inst. Zool., Acad. Sinica* **21**: 129-143.
- Murphy RW, JW Sites, DG Buth, CH Haufler. 1990. Proteins. I: Isozyme electrophoresis. *In Molecular systematics*, eds. DM Hills, C Moritz. Massachusetts: Sinauer, pp. 45-126.
- Ozaki H, Y Fujio. 1985. Genetic differentiation in geographical populations of the Pacific oyster (*Crassostrea gigas*) around Japan. *Tohoku J. Agricul. Res.* **36**: 49-61.
- Singh SM, RH Green. 1984. Excess of allozyme homozygosity in marine molluscs and its possible biological significance. *Malacologia* **25**: 569-581.
- Singh SM, E Zouros. 1978. Genetic variation associated with growth rate in the American oyster (*Crassostrea virginica*). *Evolution* **32**: 342-353.
- Sokal RR, FJ Rohlf. 1981. *Biometry*. 2nd ed. New York: W.H. Freeman and Co., pp. 859.
- Soong K, LL Liu, JL Chen, CP Chen. 1992. Regional differences of oyster (*Crassostrea gigas* Thunberg) size and shape in Taiwan. *Bull. Inst. Zool., Acad. Sinica* **31**: 111-119.
- Volckaert F, E Zouros. 1989. Allozyme and physiological variation in the scallop *Placopecten magellanicus* and a general model for the effects of heterozygosity on fitness in marine molluscs. *Mar. Biol.* **103**: 51-61.
- Zouros E, GH Pogson. 1994. The present status of the relationship between heterozygosity and heterosis. *In Genetics and evolution of aquatic organisms*, ed. AR Beaumont. London: Chapman and Hall, pp. 135-146.
- Zouros E, SM Singh, HE Miles. 1980. Growth rate in oysters: an overdominant phenotype and its possible explanations. *Evolution* **34**: 856-867.

臺灣大牡蠣族群之同功酶變異

劉莉蓮¹ 宋克義¹ 陳章波²

以蛋白質電泳法分析比較本省九個地區的大牡蠣其同功酶在族群間之變異，七個同功酶中只有 *Lap*、*Mpi* 及 *Pgi* 三個為多型性同功酶。異基因型個體缺少的現象 (heterozygote deficiency)，只在本省之七個採樣點發現，澎湖之二採樣點則無此現象。本省牡蠣族群間基因平均異質性 (mean heterozygosity)，為 $H=0.044\sim 0.084$ ，Nei's 遺傳距離為 $D<0.01$ ，顯示臺灣大牡蠣族群間遺傳變異很小。布袋族群中同功酶 *Lap* 異基因型個體顯著小於同基因型個體 ($p=0.03$)，但以多次測驗之調整顯著水準為標準時，則異基因型個體小於同基因型個體之現象不顯著。由本實驗結果推測，環境因子可能是造成本省養殖牡蠣族群間體型及大小差異之重要因素。

關鍵詞：遺傳變異，基因異質性。

¹ 國立中山大學海洋生物研究所

² 中央研究院動物研究所