

# Population Genetic Structures of the Jellyfish *Aurelia coerulea* Polyps along Korean Coasts and Implications as Revealed by Mitochondrial *COI*

Yoseph Seo<sup>1,§</sup> , Buhari Lawan Muhammad<sup>1,§</sup> , Jinho Chae<sup>2</sup>, and Jang-Seu Ki<sup>1,\*</sup> 

<sup>1</sup>Department of Biotechnology, Sangmyung University, Seoul 03016, South Korea. E-mail: akldtytpq12@gmail.com (Seo); buharilawan20@gmail.com (Muhammad); jinhochae@gmail.com (Chae)

<sup>2</sup>Marine Environmental Research and Information Laboratory, Gunpo 15850, South Korea. \*Correspondence: E-mail: kijos@smu.ac.kr (Ki). Tel: +82-2-2287-5449. Fax: +82-2-2287-0070.

<sup>§</sup>YS and BLM contributed equally to this work.

Received 1 April 2021 / Accepted 20 July 2021 / Published 12 November 2021  
Communicated by James D. Reimer

The moon jellyfish, *Aurelia coerulea*, is globally distributed, and its blooms have been responsible for severe environmental impacts. Benthic polyp populations are important for forming and maintaining medusa populations; however, their genetic structures are mostly unknown. Here, we analysed the genetic structure and phylogeographic pattern of *A. coerulea* polyps using the mitochondrial *COI* of 229 specimens collected from four different coastal regions of Korea. Molecular discrimination by *COI* assigned all polyps to *A. coerulea*. Population genetics revealed 53 haplotypes with high diversity and significant genetic structure, distinguishing two haplogroups (A and B) that coexist in all regions. Haplogroup A exhibited a star-like haplotype network pattern, while haplogroup B demonstrated a branched haplotype network pattern. Our results suggest that, the two haplogroups detected have existed in sympatry along Korean coasts. However, haplogroup A may have been established by a recent population expansion, while haplogroup B may have been established a long time ago. The strong genetic structure found within the polyp population of *A. coerulea* may have an effect on the moon jellyfish blooms on Korean coasts.

**Key words:** *A. coerulea*, Benthic polyp, Cytochrome *c* oxidase subunit I, Genetic diversity, Jellyfish bloom.

## BACKGROUND

Mass outbreaks of jellyfish have increased in the last several decades in near-shore regions and coastal ecosystems across the globe (Condon et al. 2012). The main causes of these increase in these jellyfish blooms are thought to be the rising of ocean temperatures (Condon et al. 2012), eutrophication (Arai 2008; Purcell et al. 2007), and the construction of marine structures (Duarte et al. 2013). The large aggregations (blooms) of jellyfish are associated with several negative socio-economic impacts in many harbours and coastal waters around the world (Mills 2001). They have caused

significant ecological and economic problems in the San Francisco Bay of Northern California (Greenberg et al. 1996), the Gulf of Mexico (Frolova and Miglietta 2020), Japan (Yasuda 2004), and South Korea (Ki et al. 2008; Yoon et al. 2018). These problems have ranged from fishery damage (Kim et al. 2012; Quiñones et al. 2013) to the clogging of seawater intakes of power plants and ships (Abdul Azis et al. 2000; Ki et al. 2008). In addition, jellyfish have also clogged the pumps used to flush the Foster City Lagoon, an artificial recreational pool, and fouled beaches (Greenberg et al. 1996).

The moon jelly *Aurelia* is one of the predominant bloom-forming jellyfish in coastal and shelf ecosystems.

Citation: Seo Y, Muhammad BL, Chae J, Ki JS. 2021. Population genetic structures of the jellyfish *Aurelia coerulea* polyps along Korean coasts and implications as revealed by Mitochondrial *COI*. Zool Stud 60:63. doi:10.6620/ZS.2021.60-63.

Within the genus, *Aurelia coerulea* von Lendenfeld 1884 is a taxon with the broadest world distribution (Dawson et al. 2005; Scorrano et al. 2017). Generally, *A. coerulea* is believed to have spread across oceanic regions from a single origin (Dawson et al. 2005; Ki et al. 2008). Thus, severe blooms of *A. coerulea* have been frequently observed worldwide (Fofonoff et al. 2018).

On Korean coasts, the occurrence of *A. coerulea* blooms dates back to 1980. Since then, blooms have been frequently reported in harbour and port areas, especially in the western and southern coastal areas, and caused huge economic losses estimated at US\$ 277.9 million per year (Kim et al. 2012). This has gained government attention, which has led to the establishment of a management policy that includes survey and removal of medusae and polyps (if necessary) of *A. coerulea* in Korean coastal waters (Yoon et al. 2018). However, despite enormous damage, current knowledge of the population's genetic structures, distribution, and source of the outbreak is lacking. Without knowledge of the polyp genetic structures, geographical ranges, and their possible link to medusae blooms, it is difficult to formulate effective bloom management strategies (Dawson et al. 2015; van Walraven et al. 2016).

Most studies on jellyfish blooms have focused on adult medusae. However, for many *Aurelia* spp., the magnitude and timing of medusae blooms are dependent on the dynamics of their benthic polyp population and the environmental factors that control it (Marques et al. 2019). In addition, the potential asexual reproductive ability of polyps is assumed to have a significant effect on the blooms of medusae (Arai 2008; Schiariti et al. 2014; Takao et al. 2014). During strobilation events, one polyp can produce ~100 ephyrae (Lucas et al. 2012); therefore, depending on the duration of the polyp stage, the asexual reproduction of the polyp can result in apparently irregular and unpredictable blooms of the medusae (Boero et al. 2008). However, although benthic populations of *Aurelia* spp. create large blooms of free-living medusae, little has been investigated so far, probably because of their small body size and cryptic lifestyle, which make their detection difficult (Bayha and Graham 2009).

Recent studies have focused on the benthic populations; the causes and consequences of benthic population dynamics, bloom events, and environmental tolerance have also received much attention (Lucas et al. 2012; Marques et al. 2019; Frolova and Miglietta 2020). However, studies on the population genetic structure, diversity, and phylogeography of the polyp population in *Aurelia* spp. are limited. van Walraven et al. (2016) were the first to use the sessile benthic stage of *A. aurita* to show high genetic diversity and differentiation in polyp populations similar to those in medusa

populations. However, since *A. coerulea* is the taxon with the broadest world distribution, causing greater socio-economic damage than any other *Aurelia* species, it is necessary to investigate its polyp population. This will help to determine the population differentiations that may exist in the benthic stage, and their potential implications for the medusa population. Understanding the genetic structure and phylogeography of the polyp population is necessary to predict when and where jellyfish blooms will occur (van Walraven et al. 2016).

The specific impacts of *A. coerulea* are poorly understood because it is easy to confuse this species with native cryptic species (Dawson and Jackobs 2001; Fofonoff et al. 2018). Species crypsis and morphological plasticity make the species difficult to distinguish (Dawson 2005; Scorrano et al. 2017). To resolve this, molecular tools have been applied to differentiate cryptogenic species, identify population origins, and assess the impacts of invasions. Among the molecular markers, mitochondrial cytochrome *c* oxidase subunit I (*COI*) has been widely used as a powerful molecular marker to determine the genetic variations and structure of populations in many organisms (Xu et al. 2011; Dong et al. 2015; Palraju et al. 2018; Choi et al. 2020). It is a protein-coding gene in the mitochondrial genome that serves as a useful genetic marker, providing valuable information for uncovering phylogeographic patterns and ecological adaptations (Ramšak et al. 2012). *COI* has been widely used for population genetic studies, especially intra-specific analysis, due to its fast evolution, high polymorphism, and easy amplification and sequencing (Hu et al. 2008; Xu et al. 2011; Palraju et al. 2018).

In the present study, we used the mitochondrial *COI* gene to study the population genetic structure and phylogeography of *A. coerulea* polyp populations. The samples were collected from four different regions that cover most of the areas inhabited by *A. coerulea* on the Korean coastal waters. The aim of this study was to explore the genetic structure and phylogeographic patterns of the benthic populations of bloom-forming jellyfish in Korean coastal regions. This allows us to resolve the distributions and patterns of genetic differentiation among the populations of *A. coerulea* and identify the importance of its benthic population's dynamics in medusae blooms.


## MATERIALS AND METHODS

### Study site and sample collection

A total of 229 polyp samples of *Aurelia coerulea* were collected in April 2017 from four different

geographical regions (R1, R2, R3, and R4) located on the western and southern coasts of Korea (Fig. 1). Within R1 (Jeongok, JG; Dangjin, DJ; Daesan, DS) and R2 (Baeksajang, BJ; Daecheon, DC; Galyeongdo, GL),

three locations were chosen for sampling. Within R3, two locations were chosen (Usuyeong, UY; Nokdong, ND), and in R4, only one location (Jaran Bay, GS) was chosen for the sampling. Detailed information regarding



**Fig. 1.** Sampling locations of the 229 polyp samples of *Aurelia coerulea* collected within four regions (R1, R2, R3, and R4) from western and southern coasts of Korea. Detailed information on the sampling locations are listed table 1.

**Table 1.** Details of the sampling locations of 229 *Aurelia coerulea* polyps collected from western and southern coasts of Korea

Region	LC	N	Sampling date	Location	GPS
R1	JG	15	2017.4.6	Jeongok port	37°11'20.7"N, 126°38'50.4"E
	DJ	19	2017.4.13	Dangjin port	36°59'12.2"N, 126°44'50.1"E
	DS	45	2017.5.2	Daesan port	37°00'58.7"N, 126°25'25.5"E
R2	BJ	17	2017.4.13	Baeksajang port	36°35'10.9"N, 126°18'56.9"E
	DC	20	2017.4.14	Daecheon port	36°19'41.7"N, 126°30'32.2"E
	GL	23	2017.4.4	Galyeongdo port	35°43'35.9"N, 126°31'46.3"E
R3	UY	3	2017.3.29	Usuyeong port	34°34'07.9"N, 126°18'32.3"E
	ND	33	2017.4.9~19	Nokdong port	34°31'25.8"N, 127°08'06.9"E
R4	GS	54	2017.3.30	Jaran Bay	34°54'59.6"N, 128°15'11.5"E

LC, location code; N, Number of sample.

the regions and sampling locations are listed in table 1.

Polyp samples were collected by scuba divers who were trained to dive for polyp sampling and distinguish polyps *in situ*; divers also sketched underwater polyp distribution and took photographs. Collection proceeded by scraping the polyps and their substrate using a filling knife, and then placing collected scrapings into a 1-L Kautex jar. Collected samples were immediately fixed with 100% ethanol and transported to the laboratory. *Aurelia coerulea* polyps were identified morphologically, and individuals were separated using tweezers under a dissecting microscope.

### DNA extraction, amplification, and sequencing

Ethanol-preserved polyps were washed individually several times with distilled water, then genomic DNA (gDNA) was extracted using the modified cetyltrimethylammonium bromide (CTAB) method (Ausubel et al. 1989).

Polymerase chain reaction (PCR) amplification of *COI* fragments was done by using the primer pair Fco04 5'-CAA CAA ACC ATA AAG ATA TAG GAA C-3' (Folmer et al. 1994; slightly modified) and Rco800 5'-TAT TGC TAT CAT AGC ATA AAC CAT-3' (newly designed in this study). In brief, 20  $\mu$ L reaction solution was prepared for PCR. The reaction solution consisted of 2  $\mu$ L template DNA, 2  $\mu$ L Ex Taq buffer, 2  $\mu$ L dNTP mix, 1  $\mu$ L forward primer, 1  $\mu$ L reverse primer, 0.2  $\mu$ L Ex Taq Polymerase (TaKaRa Bio Inc., Shiga, Japan), and 11.8  $\mu$ L distilled water. PCR reaction conditions were as follows: 94°C for 3 min, followed by 40 cycles of 94°C for 30 s, 55°C for 30 s, and 72°C for 1 min, and finally 72°C for 5 min. PCR results were confirmed on a 1% agarose gel using 60  $\mu$ L/400 mL gel MIDORI<sup>Green</sup> (Nippon Genetics Europe GmbH, Germany) as a fluorescent source. Confirmed samples were purified using the PCR Cleanup S & V Kit (Bionics, Daejeon, Korea) according to the manufacturer's instructions.

Purified PCR products were sequenced by Bionics Inc. (Seoul, Korea) using the same primer pair as used for PCR amplification. Sequencing was performed using the BigDye<sup>TM</sup> Terminator v3.1 Cycle Sequencing Kit (Thermo Fisher Scientific) and the synthesised sequences were analysed using an Applied Biosystems 3730xl DNA Analyser (Applied Biosystems, Foster City, CA). Sequencing data were compared with reference sequences available in the GenBank database. All sequences were identified as *A. coerulea* and compiled in MEGA X software (Kumar et al. 2018). *COI* sequences were translated into amino acid sequences using the coelenterate mitochondrial code in MEGA to confirm open reading frames. The consensus sequences were trimmed to 759-bp fragments

and aligned using Clustal Omega (Sievers et al. 2011). All sequences determined in this study were deposited in the GenBank database under the accession numbers MT251296-MT251346 and MT361137-MT361314. A list of all sequences used in this study is provided in table S1.

### Population and phylogenetic analyses

The assembled sequences were imported into Arlequin v.3.5.2.2 (Excoffier and Lischer 2010) to calculate the genetic diversity indices (number of haplotypes (Nh), number of polymorphic sites (Ps), haplotype diversity (h), and nucleotide diversity ( $\pi$ ). In addition, Tajima's (*D*) (Tajima 1989) and Fu's (*F<sub>s</sub>*) scores (Fu 1997) were calculated using DnaSP v.6 (Rozas et al. 2017).

To explore the phylogeographic patterns of the present *A. coerulea* haplotypes, we constructed a TCS haplotype network (Clement et al. 2002) for the 229 *COI* sequences using PopART v1.7 (Leigh and Bryant 2015). Haplotype networks were constructed using the default settings. Genetic variation among the different clusters was tested. A total of 229 *COI* sequences (759 bp) were imported into Arlequin v.3.5.2.2, where Analysis of Molecular Variance (AMOVA) was assessed, and a pairwise *F<sub>ST</sub>* matrix with 10,000 permutations was constructed. The haplotype distance matrix was graphically rendered using R-3.0 software (R Core Team 2014). Principal Component Analysis (PCA) was used to further evaluate and visualise the geographic genetic structure among the populations, and was calculated in Palaeontological Statistics Software Package (PAST) v.3.25 (Hammer et al. 2001), using the haplotype frequency of the 53 unique haplotypes (Table S2).

For phylogenetic analysis, we constructed a dataset of 53 *COI* reference sequences of the haplotypes determined in this study and 115 *COI* sequences of *A. coerulea* (*Aurelia* sp.1) retrieved from the GenBank database. (Table S3). Overall, 168 *COI* sequences (584 bp) were aligned using Sequencher v.5.1, (Gene Codes Corporation, Ann Arbor, MI), and the alignment results were trimmed to remove ambiguous regions. A phylogenetic tree was constructed using MEGA X software (Kumar et al. 2018). The tree was inferred using neighbor-joining (NJ) computed with the Kimura 2-parameter model (Kimura 1980), and taxa were clustered together with 1000 bootstrap values. The phylogenetic tree was visualised with Mega X Tree Explorer and edited with Adobe Illustrator CS6 (Adobe Systems, San Jose, CA).

## RESULTS

### Genetic diversity

In this study, a total of 229 *Aurelia coerulea* specimens were analyzed. Our BLAST search comparisons for each of the 229 *COI* sequences revealed that our sequences shared 100% of nucleotide identity with a known *COI* DNA sequence from *A. coerulea* (GenBank accession no. LK022703, formerly *Aurelia* sp.1). Our genetic diversity analysis of 229 *Aurelia coerulea* polyps identified 51 polymorphic sites (Ps), and represents 53 haplotypes (Table 2). The overall haplotype diversity ( $h$ ) and nucleotide diversity ( $\pi$ ) across the four geographic regions were estimated to be  $0.8788 \pm 0.0169$  and  $0.005532 \pm 0.003052$ , respectively. The highest haplotype diversity was calculated for R3 (0.9), while the lowest was for R1 (0.8079). The highest nucleotide diversity was also observed in R3 (0.0056471), and the lowest was observed in R2 (0.004577). Neutrality tests of the *COI* haplotypes were performed for all populations. Tajima's  $D$  and Fu's  $F_s$  values were negative for all populations, but only Fu's  $F_s$  values at R2 and R4 were statistically significant. However, the overall values of both Tajima's  $D$  and Fu's  $F_s$  showed significantly negative values ( $D = -1.484$  ( $p < 0.1$ ) and  $F_s = -25.442$  ( $p < 0.05$ ) (Table 2).

### Genetic differentiation

AMOVA analysis was carried out for the populations based on their geographical distance. The analysis showed that most of the genetic differentiation occurred within populations (83.87%,  $p < 0.001$ ), while 18.88% of the genetic differentiation occurred among populations within regions, and -2.76% occurred among regions (Table 3). The overall  $F_{ST}$  value ( $F_{ST} = 0.162$ ;  $p = 0$ ) suggested moderate but significant genetic differentiation within populations with significant gene flow. Pairwise  $F_{ST}$  analysis showed the pattern of genetic differentiation for all locational populations, and analysis showed that most of the significant genetic differentiation was between location DS in R1 and the rest of the locations (Fig. 2). The largest level of genetic differentiation was found between R1 (DS) and R2 (GL), while the smallest genetic differentiation was found between location UY in R3 and other locations, except for DS in R1. However, the  $F_{ST}$  test showed no significant genetic differentiation among the rest of the locations.

### Phylogeography and distribution of *A. coerulea* on Korean coasts

The TCS haplotype network analysis of 229 *A.*


**Table 2.** Genetic diversity indices for the *COI* of 229 polyp samples of *Aurelia coerulea* collected from four regions (R1, R2, R3, and R4) on western and southern coasts of Korea

Region	N	Nh	Ps	D	$F_s$	$h$	$\pi$
R1	79	21	27	-0.81	-5.215	$0.8079 \pm 0.0265$	$0.005316 \pm 0.002973$
R2	60	20	27	-1.284	-7.152*	$0.8554 \pm 0.0349$	$0.004577 \pm 0.002626$
R3	36	15	22	-0.654	-3.301	$0.9 \pm 0.0348$	$0.005647 \pm 0.003183$
R4	54	21	27	-1.05	-7.541*	$0.8819 \pm 0.032$	$0.00529 \pm 0.002979$
Total	229	53	51	-1.487	-25.442*	$0.8788 \pm 0.0169$	$0.005532 \pm 0.003052$


N, Number of sample; Nh, Number of haplotype; Ps, Number of polymorphic site; D, Tajima's  $D$  score;  $F_s$ , Fu's  $F_s$  score;  $h$ , Haplotype diversity;  $\pi$ , Nucleotide diversity; \*,  $p < 0.05$ .

**Table 3.** Analysis of molecular variance (AMOVA) results showing partitioning of genetic variation of (A) all populations and (B) between two haplogroups (A and B)

Source of variation	d.f.	Sum of Squares	Variance Components	% of Variation	Fixation indices	P-value
(A) All regions						
Among regions	3	37.792	-0.06	-2.78%	$F_{CT} = -0.028$	Not significant
Among locations Within regions	5	46.376	0.409	18.94%	$F_{SC} = 0.184$	$p = 0$
Within Populations	220	398.667	1.812	83.84%	$F_{ST} = 0.162$	$p = 0$
(B) Haplogroups (A and B)						
Among haplogroups	1	278.06	2.673	77.35%	$F_{ST} = 0.774$	$p = 0$
Within Populations	216	169.066	0.783	22.65%		



**Fig. 2.** Pairwise  $F_{ST}$  data based on the 53 *COI* haplotypes of *Aurelia coerulea* analyzed between nine locations, within four regions (R1, R2, R3 and R4) of Korean coasts. The abbreviations of the locations are listed in table 1.




**Fig. 3.** Haplotype (TCS) network showing the phylogeography of the 53 *COI* haplotypes of *Aurelia coerulea* collected from western and southern coasts of Korea. Each circles represent individual haplotypes and the size of the cycle is the proportion of haplotype frequency. The color of each segment indicates the geographic origin of that fraction of the samples. Each line between haplotypes represents a 1-nucleotide mutational change. Small black dots indicate unsampled haplotypes.

*coerulea* polyps revealed the phylogenetic similarities and geographical distribution of *COI* haplotypes in Korean coastal waters. Overall, the 53 haplotypes were divided into two distinct haplogroups (A and B) (Fig. 3). Haplogroup A was formed by 19 haplotypes; all the haplotypes in the group were linked to the most frequent haplotype (H04), showing a star-like phylogeny consistent with a pattern of population expansion. In contrast, haplogroup B comprised 24 haplotypes that formed a branched haplotype network. The two haplogroups were separated by five mutational steps, and there were 10 intermediate haplotypes that did not belong to either group (H02, H15, H17, H32, H33, H34, H38, H39, H41, and H45). Overall, the most frequent haplotype was H04 from haplogroup A (71 individuals), and this haplotype was shared among all four regions. Within haplogroup B, H09 was the most frequent haplotype. In total, there were 13 haplotypes that shared more than one region (Fig. 3), and the rest were unique to a single region. H26 and H06 were the most frequent unique haplotypes found in haplogroup A and B, respectively. AMOVA analysis based on the two haplogroups (A and B) yielded a significant genetic variation, with 77.35% of the genetic differentiation

occurring among populations and 22.65% occurring within populations. The  $F_{ST}$  value was 0.774, highlighting a gene flow restriction between the two haplogroups (Table 3).

The 53 haplotypes were distributed across the four regions of Korean coastal waters. The haplotypes belonging to the two haplogroups coexisted in all four regions, despite significant genetic variation between the two haplogroups (Fig. 4). As mentioned above, haplotype H04 was the most frequently found haplotype in all four regions. However, in R1, H09 and H06 were codominant with H04 (Fig. 4). Analysis of the PCA was carried out to summarize the genetic diversity on a geographical basis. The analysis revealed the distribution of all 53 haplotypes in the four regions (Fig. 5). Each green arrow represents the geographical locations, with angles between arrows indicating the significance of the correlation and the distance between the dots in the diagram approximating the dissimilarity of the haplotypes in the four regions. Most of the haplotypes (Black dots) are less frequent and have low variation in the regions. Haplotypes H09 and H06 in R1 and H26 in R2 showed the greatest variation, while H04 is the most frequent.



**Fig. 4.** Distribution of the *COI* haplotypes of *Aurelia coerulea* in four regions (R1, R2, R3 and R4) in the Korean coastal waters. The small circles represent the proportion of haplogroup A, haplogroup B and the intermediate haplotypes in each region. The big circles represent the proportion of the haplotypes found in each region, and the number in the middle indicates the total number of individuals sampled in each region. Each color in the circles represents the frequencies of the dominant haplotype. Others are the total frequencies of the haplotypes that are not shown in the respective circles.




## Phylogenetic analysis

The present phylogenetic tree was constructed based on the neighbor-joining (NJ) method using the *COI* gene sequences of our 53 haplotypes and 115 *COI* sequences of the global *A. coerulea* population retrieved from the GenBank database (ESM 1, Table S2 and S3). The tree showed that all global *A. coerulea* populations were divided into two main clades that were similar to the TCS haplotype network of the Korean populations. All haplotypes in haplogroup A from the TCS network clustered in Clade I (Fig. S1A), while all the haplotypes in haplogroup B clustered in Clade II (Fig. S1B). Haplotypes from Japan, China, Europe, and Australia nested in both clades. However, most of the European and Australian haplotypes nested in Clade I. In addition, American haplotypes clustered only in Clade I. The intermediate haplotypes in TCS network were indistinguishable and clustered in Clade I of the phylogenetic tree.

## DISCUSSION

### Genetic structure

In the present study, we found high genetic diversity in all regions, especially in R3 and R4. Overall, high genetic diversity was found in Korean coastal waters, which was similar to a previous finding



**Fig. 5.** PCA plot showing the distribution of *Aurelia coerulea* haplotypes in the four regions (R1, R2, R3, and R4) of Korean coastal waters. The black dots are the haplotypes that are less frequent and have less variation in the four region. The coloured dots are haplotypes that are more frequent and have more variation between the four regions. The green arrows represent parameters pointing in the direction of the four geographical locations (R1, R2, R3, and R4).

in Japanese populations ( $h = 0.87 \pm 0.06$ ;  $\pi = 0.0063 \pm 0.0037$ ) (Dawson et al. 2005). The high genetic diversity found in this study was expected because *Aurelia coerulea* is believed to be a native species in the western North Pacific (Dawson et al. 2005). In addition, Fofonoff et al. (2018) mentioned that the species is most likely a native from southern Korea to northern Japan and possibly China. Usually, high genetic diversity is observed in the native populations and is favoured by a large stable population size and positive demographic balance (Grant and Bowen 1998; Micheli et al. 2015), which improves the resistance and resilience of populations to environmental changes (Roger et al. 2012). The genetic data presented here support the idea that *A. coerulea* is likely native to the western North Pacific, as suggested by Dawson et al. (2005). However, the negative and significant values shown by Tajima's  $D$  and Fu's  $F_s$  tests indicated that there has been a recent population expansion and an increase in effective population size in Korean coastal regions.

The AMOVA analysis based on all *A. coerulea* populations showed that most of the genetic differentiation occurred within populations. This implies that there was no restricted gene flow between the four regions (except between R1 and R2) and shows a sympatric distribution of the two haplogroups. Restricted gene flow is caused primarily by geographical barriers and low dispersal ability of species (Liu et al. 2019). Pairwise  $F_{ST}$  analysis showed that most of the genetic differentiation occurred between region R1, specifically DS, and the rest of the regions. This is apparently due to haplotypes H06 and H09, which were mainly found in DS (Fig. S2). Similarly, haplotype H26 was restricted to R2, resulting in unequal gene flow and high genetic differentiation between R1 and R2 despite a short geographic distance. This is clearly shown in the PCA analysis. Usually, genetic differentiation between more distant populations is higher due to substantial differences in allele frequencies (Liu et al. 2019), which could be interpreted as isolation-by-distance or allopatric speciation (Coyne and Orr 2004; Zhan et al. 2009). However, in this case, it could not be isolation-by-distance; rather, R1 and R2 displayed a divergent pattern similar to parapatric speciation (Butlin et al. 2008). That is, there was restricted gene flow between the diverging populations. The reason for this unique differentiation between R1 and R2 may be due to hydrodynamic circulation, habitat fragmentation formed by marine gyres and currents, or even anthropogenic activities (Otto et al. 1990; Launey et al. 2002; Zhan et al. 2009; Dong et al. 2015; Liu et al. 2019).



## Phylogeography and intra-specific variation in *A. coerulea*

Our TCS haplotype network analysis revealed a well-resolved phylogeography of *A. coerulea* in Korean coastal waters. The 229 polyp samples collected were clearly divided into two major haplogroups (A and B) separated by five mutational steps with intermediate haplotypes. AMOVA constructed based on these two distinct haplogroups showed high genetic differentiation among populations with a high  $F_{ST}$  value of 0.77353. This clearly indicates high genetic variation and restricted gene flow between the two groups. The haplogroups displayed different patterns, with haplogroup A having a star-like haplotype network, providing support for a recent population expansion, as described by Avise (2000) and Froufe et al. (2016). Haplogroup B showed a branched haplotype network, which is thought to be the nature of a stable and older clade (Ferrerri et al. 2011).

Interestingly, the haplotypes of the two haplogroups coexisted in all regions in Korean coastal waters, and the coexistence of these divergent haplogroups in the same geographical locations suggests sympatric development (Avise 2000). Population subdivision and intraspecific phylogeographic structures are commonly observed in jellyfish (e.g., Holland et al. 2004; Dawson 2005; Ramšak et al. 2012; Lee et al. 2013) and other organisms, including bluegill sunfish (*Lepomis macrochirus*; Avise et al. 1984), rock slaters (*Ligia occidentalis*; Markow and Pfeiler 2010), and freshwater mussels (Potomida, Unionida; Froufe et al. 2016). However, these are all examples of allopatric differentiation, the most commonly encountered situation. It has been rare to observe large mtDNA differences within the same species collected at any given geographic site (Avise et al. 1987). Though this large mtDNA difference was recently reported in the jellyfish *Pelagia noctiluca* (Stopar et al. 2010; Glynn et al. 2016), we did not find anything like this reported in *A. coerulea*.

Yi and Kim (2020) found similar significant genetic differentiation among ascidian *Ciona savignyi* populations along Korean coasts. They suggested that a possible northward post-glacial colonisation scenario in the western North Pacific around 15 thousand years ago left evidence of the expansion and differentiated genetic structure of *C. savignyi* populations and other coastal marine organisms in Korea (Yi and Kim 2020). We did not estimate molecular clock calibration for scyphozoans in the present study, because a reliable molecular clock calibration for scyphozoans is not available (Holland et al. 2004); therefore, the estimated separation time could not be determined. However,

with high genetic variation and restricted gene flow found between the two haplogroups in this study, the above hypothesis gives a possible explanation for the *COI* variation found in *A. coerulea*. Based on this and inferences made by several previous studies, Korean coastal regions may be considered a zone of secondary admixture between allopatrically evolved populations (Avise et al. 1987; Avise et al. 1998; Avise 2000; Hewitt 2000; Beebe and Rowe 2004; Provan and Bennett 2008).

In addition, for the secondary contact and expansion event, haplogroup B, with its diverse network topology, might be the older clade that settled a long time ago in Korean coastal waters. However, the star-like phylogeny of haplogroup A is generally interpreted as indicative of a population that recently expanded (Slatkin and Hudson 1991; Avise 2000; Froufe et al. 2016). This is supported by the overall neutrality test from this study. Therefore, there is strong evidence for past population expansion through introductions by anthropogenic means or natural dispersal, shaping the genetic structure of *A. coerulera* in Korean coastal waters.

## Global distribution of *A. coerulea*

As mentioned above, the moon jellyfish *A. coerulea* has the broadest geographic range of all species in the genus *Aurelia* to date (Ki et al. 2008; Scorrano et al. 2017). The species was suggested to be native to the western North Pacific (Japan, Korea, and China) and dispersed globally from Japan by anthropogenic means (Dawson et al. 2005). This is supported by the present phylogenetic analysis based on the global *A. coerulea* population, which showed that haplotypes in haplogroup A and B are widely distributed globally, and the most frequent haplotype was clustered with Chinese and Japanese haplotypes. Due to the overlapping distribution of the two haplogroups in Korean coastal regions, it is difficult to determine the geographical origin of the two haplogroups. However, the present data highlight the possibility that, after the speciation events, haplogroup B became largely concentrated in the Bohai Sea and the Yellow Sea, while haplogroup A may have concentrated in Japanese coasts and dispersed naturally, mixing locally in the western North Pacific. However, dispersal beyond this region to other global locations occurs through anthropogenic means (Dawson et al. 2005). The presence of shared haplotypes among distant locations highlights recent connections through human-mediated dispersal (Lejeusne et al. 2014).

## Polyp populations and possible implications for the medusa bloom

Previous studies on population genetics have mostly investigated the medusa population to explore the genetic structure and diversity of *Aurelia* spp. (Dawson et al. 2005; Ramšak et al. 2012). However, medusa blooms depend on the dynamics of benthic polyp populations and environmental factors that control them (Arai 2008; Frolova and Miglietta 2020). van Walraven et al. (2016) were the first to report high genetic diversity and differentiation in *A. aurita* polyp populations, similar to medusa populations. Similarly, in the present study, we used benthic polyp samples of *A. coerulea* to explore their genetic structure and diversity in an attempt to identify the source of the medusa blooms. We found higher genetic diversity and differentiation than in our previous analysis of medusa populations in Korean coastal waters (Ki et al. 2008). The most frequent haplotype from haplogroup A (H04) in the polyp populations was also found to be the most frequent among the medusa populations (Fig. S3). This haplotype seems to be the common ancestral-like haplotype in haplogroup A, with all the haplotypes in the group connected to it by one or two mutational steps. The success of this haplotype (H04) might be due to good adaptation to environmental conditions for the transition from polyp to ephyrae, and eventually to medusa (Marques et al. 2019; Frolova and Miglietta 2020). Dawson et al. (2015) showed how genetically different populations of jellyfish of the same species respond differently to changing environmental conditions. This could result in differences in factors related to medusa blooms, such as timing and magnitude. In the present study, environmental factors were not measured; therefore, it is difficult to link the polyp population to medusa bloom. However, the significant genetic differentiation found in the polyp population which is similar to the medusa population, highlights the possible link between polyp population (Haplogroup A) and the medusa bloom, in the water column of Korean coastal waters. Further studies should be carried out to thoroughly examine how the two different haplogroups will respond to environmental changes.

## CONCLUSIONS

In the present study, high genetic diversity with a significant genetic structure within *A. coerulea* polyp populations was found. The *A. coerulea* population was divided into two haplogroups (A and B). Based on the relevant data and evidence presented in this study, we

conclude that: (1) Although strong inferences cannot be made about specific causes of the *COI* variations found in *A. coerulea*, a likely explanation for the divergence is that the current populations consist of separate lineages that have evolved in allopatry. However, additional genetic markers would provide higher-resolution information on population connectivity that would reflect the early stages of speciation. (2) The star-like haplotype network of haplogroup A suggests multiple expansion events. Therefore, haplogroup A might be established by recent population expansion through anthropogenic means or natural dispersal, as supported by phylogenetic analysis and neutrality tests. In contrast, haplogroup B might be the older clade that was established a long time ago in Korean coastal waters. (3) Genetic differentiation of the polyp population may have an effect on medusa blooms. H04 from haplogroup A matched well with the haplotype of the bloom-forming jellyfish medusa. Thus, haplotype H04 might be responsible for *A. coerulea* blooms in Korean coastal regions. However, environmental factors should be thoroughly examined to determine how the two different haplogroups respond to environmental changes.

Finally, our results reveal a strong genetic structure of the *A. coerulea* polyp population in Korean coastal waters. This could be used as reference data for future studies to resolve the global phylogeography of *A. coerulea*.

**Acknowledgments:** This research was supported by a part of the project titled “Improvement of management strategies on marine disturbing ecosystems and harmful organisms (No. 20190518)” funded by the Ministry of Oceans and Fisheries, Korea.

**Competing interests:** The authors declare no conflict of interest.

**Authors' contributions:** Seo YS: conceptualization, investigation; Muhammad BL: writing original draft, writing-review & editing. Chae JH: Investigation. Ki JS: Conceptualization, Writing - review & editing.

**Availability of data and materials:** All the data presented in this study are available in the supporting information (Supplementary materials: Figs. S1 and 2). All sequences determined were deposited in the GenBank database under the accession numbers MT251296-MT251346 and MT361137-MT361314.

**Consent for publication:** Not applicable.

**Ethics approval consent to participate:** Not applicable.

## REFERENCES

- Abdul Azis PK, Al-Tisan I, Al-Daili M, Green TN, Dalvi AGI, Javeed MA. 2000. Effects of environment on source water for desalination plants on the eastern coast of Saudi Arabia. *Desalination* **132**:29–40. doi:10.1016/S0011-9164(00)00132-6.
- Arai MN. 2008. The potential importance of podocysts to the formation of scyphozoan blooms: a review. In: Pitt KA, Purcell JE (ed) *Jellyfish blooms: Causes, consequences and recent advances*. Springer, Dordrecht, pp. 241–246. doi:10.1007/978-1-4020-9749-2\_16.
- Ausubel FM, Brent R, Kingston RF, Moore DD, Seidman JG, Smith JA, Struhl K. 1989. *Current Protocols in Molecular Biology*. Wiley and Sons, New York, USA.
- Avise JC. 2000. *Phylogeography: the history and formation of species*. Harvard University Press, Cambridge, UK.
- Avise JC, Arnold J, Ball RM, Bermingham E, Lamb T, Neigel JE, Reeb CA, Saunders NC. 1987. Intraspecific Phylogeography: The mitochondrial DNA bridge between population genetics and systematics. *Annu Rev Ecol Evol Syst* **18**:489–522. doi:10.1146/ANNUREV.ES.18.110187.002421.
- Avise JC, Bermingham E, Kessler LG, Saunders NC. 1984. Characterization of mitochondrial DNA variability in a hybrid swarm between subspecies of bluegill sunfish (*Lepomis macrochirus*). *Evolution* **38**:931–941. doi:10.1111/j.1558-5646.1984.tb00364.x.
- Avise JC, Walker D, Johns GC. 1998. Speciation durations and Pleistocene effects on vertebrate phylogeography. *Proc Royal Soc B* **265**:1707–1712. doi:10.1098/rspb.1998.0492.
- Bayha KM, Graham WM. 2009. A new Taqman<sup>®</sup> PCR-based method for the detection and identification of scyphozoan jellyfish polyps. In: Pitt KA, Purcell JE (ed) *Jellyfish Blooms: Causes, Consequences, and Recent Advances*. Springer, Dordrecht, pp. 217–228. doi:10.1007/978-1-4020-9749-2\_14.
- Beebe TJ, Rowe G. 2004. *An introduction to molecular ecology*. Oxford University Press, Oxford, pp. 167–176.
- Boero F, Bouillon J, Gravili C, Miglietta MP, Parsons T, Piraino S. 2008. Gelatinous plankton: irregularities rule the world (sometimes). *Mar Ecol Prog Ser* **356**:299–310. doi:10.3354/meps07368.
- Butlin RK, Galindo J, Grahame JW. 2008. Sympatric, parapatric or allopatric: the most important way to classify speciation?. *Philos Trans R Soc Lond B Biol Sci* **363**:2997–3007.
- Choi EH, Kim G, Cha SH, Lee JS, Ryu SH, Suk HY, Lee YS, Baek YS, Hwang UW. 2020. Molecular phylogenetic, population genetic and demographic studies of *Nodularia douglasiae* and *Nodularia breviconcha* based on *COI* and *16S rRNA*. *Sci Rep* **10**:1–14. doi:10.1038/s41598-020-72015-5.
- Clement M, Snell Q, Walke P, Posada D, Crandall K. 2002. TCS: estimating gene genealogies. In: *Proceedings 16th International Parallel and Distributed Processing Symposium*, **2**:0184. doi:10.1109/IPDPS.2002.1016585.
- Condon RH, Graham WM, Duarte CM, Pitt KA, Lucas CH, Haddock SH, Sutherland KR, Robinson KL, Dawson MN, Decker MB, Purcell JE, Malej A, Mianzan H, Uye S, Gelcich S, Madin PL, Mills CE. 2012. Questioning the rise of gelatinous zooplankton in the world's oceans. *BioScience* **62**:160–169. doi:10.1525/bio.2012.62.2.9.
- Coyne JA, Orr HA. 2004. *Speciation*. Sinauer Associates, Sunderland, pp. 1–545.
- Dawson MN. 2005. Incipient speciation of *Catostylus mosaicus* (Scyphozoa, Rhizostomeae, Catostylidae), comparative phylogeography and biogeography in south-east Australia. *J Biogeogr* **32**:515–533. doi:10.1111/j.1365-2699.2004.01193.x.
- Dawson MN, Cieciel K, Decker MB, Hays GC, Lucas CH, Pitt KA. 2015. Population-level perspectives on global change: genetic and demographic analyses indicate various scales, timing, and causes of scyphozoan jellyfish blooms. *Biol Invasions* **17**:851–867. doi:10.1007/s10530-014-0732-z.
- Dawson MN, Gupta AS, England MH. 2005. Coupled biophysical global ocean model and molecular genetic analyses identify multiple introductions of cryptogenic species. *Proc Natl Acad Sci USA* **102**:11968–11973. doi:10.1073/pnas.0503811102.
- Dawson MN, Jacobs DK. 2001. Molecular evidence for cryptic species of *Aurelia aurita* (Cnidaria, Scyphozoa). *Biol Bull* **200**:92–96. doi:10.2307/1543089.
- Dong Z, Liu Z, Liu D. 2015. Genetic characterization of the scyphozoan jellyfish *Aurelia* spp. in Chinese coastal waters using mitochondrial markers. *Biochem Syst Ecol* **60**:15–23. doi:10.1016/j.bse.2015.02.018.
- Duarte CM, Pitt KA, Lucas CH, Purcell JE, Uye SI, Robinson K, Brotz L, Decker MB, Sutherland KR, Malej A, Madin L. 2013. “Is global ocean sprawl a cause of jellyfish blooms?”. *Front Ecol Environ* **11**:91–97. doi:10.1890/110246.
- Excoffier L, Lischer HE. 2010. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* **10**:564–567. doi:10.1111/j.1755-0998.2010.02847.x.
- Ferreri M, Qu W, Han B. 2011. Phylogenetic networks: A tool to display character conflict and demographic history. *Afr J Biotechnol* **10**:12799–12803. doi:10.5897/AJB11.010.
- Fofonoff PW, Ruiz GM, Steves B, Simkanin C, Carlton JT. 2018. National Exotic Marine and Estuarine Species Information System. Available at <http://invasions.si.edu/nemesis/>. Accessed 5 Jan. 2021.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase I subunit from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* **3**:294–299.
- Frolova A, Miglietta MP. 2020. Insights on bloom forming jellyfish (Class: Scyphozoa) in the Gulf of Mexico: environmental tolerance ranges and limits suggest differences in habitat preference and resistance to climate change among congeners. *Front Mar Sci* **7**:93. doi:10.3389/fmars.2020.00093.
- Froufe E, Prié V, Faria J, Ghamizi M, Gonçalves DV, Gürlek ME, Sousa R. 2016. Phylogeny, phylogeography, and evolution in the Mediterranean region: news from a freshwater mussel (*Potomida*, Unionida). *Mol Phylogenet Evol* **100**:322–332. doi:10.1016/j.ympev.2016.04.030.
- Fu YX. 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics* **147**:915–925. doi:10.1093/genetics/147.2.915.
- Glynn F, Houghton JD, Bastian T, Doyle TK, Fuentes V, Lilley MK, Provan J. 2016. High-resolution genetic analysis reveals extensive gene flow within the jellyfish *Pelagia noctiluca* (Scyphozoa) in the North Atlantic and Mediterranean Sea. *Biol J Linn Soc* **117**:252–263. doi:10.1111/bij.12654.
- Grant WAS, Bowen BW. 1998. Shallow population histories in deep evolutionary lineages of marine fishes: insights from sardines and anchovies and lessons for conservation. *J Hered* **89**:415–426. doi:10.1093/jhered/89.5.415.
- Greenberg N, Garthwaite RL, Potts DC. 1996. Allozyme and morphological evidence for a newly introduced species of *Aurelia* in San Francisco Bay, California. *Mar Biol* **125**:401–410. doi:10.1007/BF00346320.
- Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: paleontological statistics software package for education and data analysis. *Palaeontol Electron* **4**:9.

- Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* **405**:907–913. doi:10.1038/35016000.
- Holland BS, Dawson MN, Crow GL, Hofmann DK. 2004. Global phylogeography of *Cassiopea* (Scyphozoa: Rhizostomeae): molecular evidence for cryptic species and multiple invasions of the Hawaiian Islands. *Mar Biol* **145**:1119–1128. doi:10.1007/s00227-004-1409-4.
- Hu J, Zhang JL, Nardi F, Zhang RJ. 2008. Population genetic structure of the melon fly, *Bactrocera cucurbitae* (Diptera: Tephritidae), from China and Southeast Asia. *Genetica* **134**:319. doi:10.1007/s10709-007-9239-1.
- Ki JS, Hwang DS, Shin K, Yoon WD, Lim D, Kang YS, Lee Y, Lee JS. 2008. Recent moon jelly (*Aurelia* sp.1) blooms in Korean coastal waters suggest global expansion: examples inferred from mitochondrial *COI* and nuclear ITS–5.8S rDNA sequences. *ICES J Mar Sci* **65**:443–452. doi:10.1093/icesjms/fsn018.
- Kim DH, Seo JN, Yoon WD, Suh YS. 2012. Estimating the economic damage caused by jellyfish to fisheries in Korea. *Fish Sci* **78**:1147–1152. doi:10.1007/s12562-012-0533-1.
- Kimura M. 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* **16**:111–120. doi:10.1007/BF01731581.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* **35**:1547–1549. doi:10.1093/molbev/msy096.
- Launey S, Leduc C, Boudry P, Bonhomme F, Naciri-Graven Y. 2002. Geographic structure in the European flat oyster (*Ostrea edulis* L.) as revealed by microsatellite polymorphism. *J Hered* **93**:331–351. doi:10.1093/jhered/93.5.331.
- Leigh JW, Bryant D. 2015. POPART: full-feature software for haplotype network construction. *Methods Ecol Evol* **6**:1110–1116. doi:10.1111/2041-210X.12410.
- Lejeune C, Saunier A, Petit N, Bégue M, Otani M, Carlton JT, Green AJ. 2014. High genetic diversity and absence of founder effects in a worldwide aquatic invader. *Sci Rep* **4**:1–9. doi:10.1038/srep05808.
- Liu Y, Dietrich CH, Wei C. 2019. Genetic divergence, population differentiation and phylogeography of the cicada *Subsalsaltria yangi* based on molecular and acoustic data: an example of the early stage of speciation?. *BMC Evol Biol* **19**:1–17. doi:10.1186/s12862-018-1317-8.
- Lucas CH, Graham WM, Widmer C. 2012. Jellyfish life histories: role of polyps in forming and maintaining scyphomedusa populations. *Adv Mar Biol* **63**:133–196. doi:10.1016/B978-0-12-394282-1.00003-X.
- Markow TA, Pfeiler E. 2010. Mitochondrial DNA evidence for deep genetic divergences in allopatric populations of the rocky intertidal isopod *Ligia occidentalis* from the eastern Pacific. *Mol Phylogenetics Evol* **56**:468–473. doi:10.1016/j.ympev.2009.12.002.
- Marques R, Darnaude AM, Schiariti A, Tremblay Y, Molinero JC, Soriano S, Hately E, Colantoni S, Bonnet D. 2019. Dynamics and asexual reproduction of the jellyfish *Aurelia coerulea* benthic life stage in the Thau lagoon (northwestern Mediterranean). *Mar Biol* **166**:74. doi:10.1007/s00227-019-3522-4.
- Micheli C, D'Esposito D, Belmonte A, Peirano A, Valiante LM, Procaccini G. 2015. Genetic diversity and structure in two protected *Posidonia oceanica* meadows. *Mar Environ Res* **109**:124–131. doi:10.1016/j.marenvres.2015.06.016.
- Mills CE. 2001. Jellyfish blooms: are populations increasing globally in response to changing ocean conditions? *Hydrobiologia* **451**:55–68. doi:10.1023/A:1011888006302.
- Otto L, Zimmerman JTF, Furnes GK, Mork M, Saetre R, Becker G. 1990. Review of the physical oceanography of the North Sea. *Neth J Sea Res* **26**:161–238. doi:10.1016/0077-7579(90)90091-T.
- Palraju M, Paulchamy R, Sundaram J. 2018. Population genetic structure and molecular diversity of *Leucinodes orbonalis* based on mitochondrial *COI* gene sequences. *Mitochondrial DNA Part A* **29**:1231–1239. doi:10.1080/24701394.2018.1436169.
- Provan J, Bennett KD. 2008. Phylogeographic insights into cryptic glacial refugia. *Trends Ecol Evol* **23**:564–571. doi:10.1016/j.tree.2008.06.010.
- Purcell JE, Uye SI, Lo WT. 2007. Anthropogenic causes of jellyfish blooms and their direct consequences for humans: a review. *Mar Ecol Prog Ser* **350**:153–174. doi:10.3354/meps07093.
- Quiñones J, Monroy A, Acha EM, Mianzan H. 2013. Jellyfish bycatch diminishes profit in an anchovy fishery off Peru. *Fish Res* **139**:47–50. doi:10.1016/j.fishres.2012.04.014.
- R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at <http://www.R-project.org/>. Accessed 5 Jan. 2021.
- Ramšak A, Stopar K, Malej A. 2012. Comparative phylogeography of meroplanktonic species, *Aurelia* spp. and *Rhizostoma pulmo* (Cnidaria: Scyphozoa) in European Seas. In: Pitt KA, Lucas CH (ed) *Jellyfish blooms*. Springer, Dordrecht, pp. 69–80. doi:10.1007/978-94-007-5316-7\_6.
- Roger F, Godhe A, Gamfeldt L. 2012. Genetic diversity and ecosystem functioning in the face of multiple stressors. *PLoS ONE* **7**:e45007. doi:10.1371/journal.pone.0045007.
- Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos Onsins SE, Sánchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism analysis of large data sets. *Mol Biol Evol* **34**:3299–3302. doi:10.1093/molbev/msx248.
- Schiariti A, Morandini AC, Jarms G, von Glehn Paes R, Franke S, Mianzan H. 2014. Asexual reproduction strategies and blooming potential in Scyphozoa. *Mar Ecol Prog Ser* **510**:241–253. doi:10.3354/meps10798.
- Scorrano S, Aglieri G, Boero F, Dawson MN, Piraino S. 2017. Unmasking *Aurelia* Shaik, species in the Mediterranean Sea: an integrative morphometric and molecular approach. *Zool J Linn Soc* **180**:243–267. doi:10.1111/zoj.12494.
- Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* **7**:539. doi:10.1038/msb.2011.75.
- Slatkin M, Hudson RR. 1991. Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations. *Genetics* **129**:555–562.
- Stopar K, Ramšak A, Trontelj P, Malej A. 2010. Lack of genetic structure in the jellyfish *Pelagia noctiluca* (Cnidaria: Scyphozoa: Semaestomeae) across European seas. *Mol Phylogenetic Evol* **57**:417–428. doi:10.1016/j.ympev.2010.07.004.
- Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**:585–595. doi:10.1093/genetics/123.3.585.
- Takao M, Okawachi H, Uye S. 2014. Natural predators of polyps of *Aurelia aurita* s.l. (Cnidaria: Scyphozoa: Semaestomeae) and their predation rates. *Plankton Benthos Res* **9**:105–113. doi:10.3800/pbr.9.105.
- van Walraven L, Driessen F, van Bleijswijk J, Bol A, Luttikhuisen PC, Coolen JWP, Bos OG, Gittenberger A, Schrieken N, Langenberg VT, van der Veer HW. 2016. Where are the polyps? Molecular identification, distribution and population differentiation of *Aurelia aurita* jellyfish polyps in the southern North Sea area. *Mar Biol* **163**:172. doi:10.1007/s00227-016-2945-4.
- Xu ZH, Chen JL, Cheng DF, Liu Y, Frederic F. 2011. Genetic

- variation among the geographic population of the grain aphid, *Sitobiona venae* (Hemiptera: Aphididae) in China inferred from mitochondrial *COI* gene sequence. *Agric Sci China* **10**:1041–1048. doi:10.1016/S1671-2927(11)60092-8.
- Yasuda T. 2004. Massive blooms of the giant medusa in Japanese waters from late summer to winter in 2002. *Bull Plankton Soc Japan* **51**:34–37. (in Japanese).
- Yi CH, Kim W. 2020. Assessing cryptic Invasion state: fine-scale genetic analysis of *Ciona savignyi* population in putative native habitat of the Korean coast. *Ocean Sci J* **55**:99–113. doi:10.1007/s12601-019-0041-7.
- Yoon W, Chae J, Koh BS, Han C. 2018. Polyp removal of a bloom forming jellyfish, *Aurelia coerulea*, in Korean waters and its value evaluation. *Ocean Sci J* **53**:499–507. doi:10.1007/s12601-018-0015-1.
- Zhan A, Hu J, Hu X, Zhou Z, Hui M, Wang S, Bao Z. 2009. Fine-scale population genetic structure of Zhikong scallop (*Chlamys farreri*): do local marine currents drive geographical differentiation? *Mar Biotechnol* **11**:223–235. doi:10.1007/s10126-008-9138-1.

## Supplementary materials

**Fig. S1.** Phylogenetic analysis showing the global distribution of *Aurelia coerulea* collected from Korean coasts. The tree was constructed based on the *A. coerulea COI* sequences determined in this study (marked in red) and the global *A. coerulea* (*Aurelia* sp.1) retrieved from the GenBank database. All the *A. coerulea* populations were divided into two main clades: Clade I (a) and Clade II (b). The tree was inferred using Neighbor-joining (NJ) computed with Kimura 2-parameter model (Kimura 1980), the taxa were clustered together on the 1000 bootstrap value and the numbers at the nodes (> 50 are shown) are bootstrap values (percentage). (download)

**Fig. S2.** Haplotype (TCS) network showing the phylogeography of the *COI* haplotypes of *A. coerulea* sampled from three locations (Daesan port DS, Jeongok port JG, and Dangjin port DJ) in Region 1 (R1) from western coasts of Korea. Each circle represents an individual haplotype and the size of the circle is the proportion of haplotype frequency. Each line between haplotypes represents 1-nucleotide mutational change. Small black dots indicate unsampled haplotypes. (download)

**Fig. S3.** TCS haplotype network showing the phylogeography of the *COI* haplotypes of *A. coerulea* from the medusa samples randomly taken from Korean coastal waters. Each circle represents an individual haplotype and the size of the circle is the proportion of haplotype frequency. Each line between haplotypes represents a 1-nucleotide mutational change. Small black dots indicate unsampled haplotypes. (download)

**Table S1.** List of all the *A. coerulea COI* sequences determined in this study. (download)

**Table S2.** Frequency and distribution of the 53 unique haplotypes detected in four different regions (R1, R2, R3, and R4) of South Korean coastal waters. (download)

**Table S3.** List of *COI* sequences of *A. coerulea* (*Aurelia* sp.1) retrieved from the GenBank database and used in the phylogenetic analysis. (download)