

# Use of DNA Barcode Sequences for Distinguishing the Three Species in the Arctic Warbler (*Phylloscopus borealis*) Species Complex

Shun-Jen Cheng<sup>1</sup>  and Yu-Cheng Hsu<sup>1,\*</sup> 

<sup>1</sup>Department of Natural Resource and Environmental Studies, National Dong Hwa University, Hualien 97401, Taiwan.

\*Correspondence: E-mail: ycsheu@gms.ndhu.edu.tw (Hsu)

E-mail: csjprinia@gmail.com (Cheng)

Received 21 December 2023 / Accepted 26 June 2024 / Published 12 November 2024

Communicated by Chih-Ming Hung

The Arctic warbler (*Phylloscopus borealis*) species complex is commonly present in the Palearctic region. By 2014, the three bird subspecies were split into three species, Arctic warbler (*P. borealis*), Japanese leaf warbler (*P. xanthodryas*), and Kamchatka leaf warbler (*P. examinandus*), based on different breeding areas and distinct vocalizations. However, their similar coloration and body size make it difficult to distinguish these species in the nonbreeding season. Taiwan is located in the potential migration routes of the Arctic warbler species complex; however, no confirmed record of *P. xanthodryas* and *P. examinandus* exists. In this study, we compared the mitochondrial cytochrome c oxidase subunit 1 (CO1) sequences of samples from breeding sites during the breeding season and confirmed that the three species could be distinguished based on CO1 gene sequences. We also examined the species of the Arctic warbler species complex samples collected from eastern Taiwan. For the first time, we confirmed that all three species visited Taiwan during migration season. In the Taiwanese samples, no clear distinction could be made between species based on plumage coloration and size, indicating that these traits are not reliable for species identification. Reassessment of the CO1 gene sequences of the three species deposited in the Barcode of Life Data System revealed that the taxonomic status needs to be updated for 31.8% of the samples.

**Key words:** Arctic warbler species complex, DNA barcode, *Phylloscopus borealis*, *Phylloscopus xanthodryas*, *Phylloscopus examinandus*

## BACKGROUND

Understanding the species list of a region is the first step in biodiversity conservation. A complete species list can be used as a basis for assessing species survival, migration, and threats; this in turn contributes to biodiversity conservation and natural resource management (Mace 2004; Kirby et al. 2008; Rutt et al. 2019; Cazalis et al. 2020; Exantus and Cézilly 2023). *Phylloscopus borealis* is a bird species belonging to the

family Phylloscopidae. It is widely distributed in the Palearctic region. Historically, three subspecies have been identified: *P. b. borealis* (Blasius, 1858), *P. b. xanthodryas* (Swinhoe, 1863), and *P. b. examinandus* (Stresemann, 1913) (Vaurie 1954). Because these subspecies cannot be easily distinguished based on morphology, they are collectively referred to as the Arctic warbler species complex or superspecies (Martens 2010). In addition to differences in body size (Saitoh et al. 2008), genetic differentiation has been reported

between Arctic warbler populations or subspecies. Reeves et al. (2008) found that the birds breeding on Sakhalin Island and the Kamchatka Peninsula differed from the Palearctic/Beringian birds in terms of their mitochondrial ND2 sequence. Saitoh et al. (2010) and Withrow et al. (2016) reported significant differences in mitochondrial cytochrome *b* gene sequences between the three subspecies. Based on allopatry in breeding areas, differences in vocalizations, and divergence of mitochondrial gene sequences, Alström et al. (2011) argued that reproductive isolation exists in the Arctic warbler species complex, and that this complex should be split into three distinct species. In the Clements Checklist of Birds of the World in 2014, the Arctic warbler species complex was formally split into three species: Arctic warbler (*P. borealis*), Japanese leaf warbler (*P. xanthodryas*), and Kamchatka leaf warbler (*P. examinandus*) (Clements et al. 2014).

The three species of the Arctic warbler species complex have distinct breeding ranges: The Arctic warbler breeds in northern Europe, northern Siberia, Far East Russia, Mongolia, northeastern China, and Alaska; the Japanese leaf warbler breeds between Honshu and Kyushu, Japan; and the Kamchatka leaf warbler breeds in the Kamchatka Peninsula, Sakhalin, Kuril Islands, and Hokkaido (del Hoyo et al. 2020a b; Lowther and Sharbaugh 2020). After breeding season, these species migrate southward for wintering via different routes. The Arctic warbler migrates along eastern China and eastern and southeastern Asia and winters in Borneo and Indonesia (Lowther and Sharbaugh 2020). A recent study revealed that the Alaskan population further migrates through the Philippines and winters in the Palau region (Adams et al. 2022). The Japanese leaf warbler is speculated to winter from Taiwan to Java, and the Kamchatka leaf warbler possibly migrates through Japan and South Korea and winters in the Philippines (del Hoyo et al. 2020a b). The Arctic warbler species complex can be found in Taiwan in nonbreeding season. According to Severinghaus et al. (2012), the Arctic warbler and Japanese leaf warbler winter in Taiwan. However, as per a recent checklist, the Arctic warbler is regarded as a common winter visitor in Taiwan. So far, no confirmed records of the Japanese leaf warbler and Kamchatka leaf warbler exist in Taiwan (Ding et al. 2023).

During breeding season, the three species can be distinguished by their distinct breeding areas and vocalizations (Alström et al. 2011). After breeding season, they can only be distinguished based on morphology at migratory stopovers or wintering sites. Minor differences exist in plumage coloration between the species: the Arctic warbler has the grayest upperpart and whitest underpart, the Japanese leaf warbler has

an even yellow coloration, and the Kamchatka leaf warbler has a yellowish strip from the central throat to underparts (Hsiao and Li 2017). Lowther and Sharbaugh (2020) and del Hoyo et al. (2020a b) have noted that the Arctic warbler is plain olive on top (tinged grayish and brownish) and paler and whitish below, with a grayish wash or streaks on the breast side and flanks; the Japanese leaf warbler has a bright olive green crown and upperparts, variably yellowish throat and underparts, and white or whitish belly; and the Kamchatka leaf warbler has an olive green, less bright crown and upperparts, with a duller, less bright, yellow throat and underside. In addition to plumage coloration, Saitoh et al. (2008) compared samples from breeding areas and noted differences in body size in the species complex. In general, the Arctic warbler is the smallest, whereas the Japanese leaf warbler is the largest. The body size of the Kamchatka leaf warbler differs between populations: the body size of the Kamchatka population is similar to that of the Japanese leaf warbler, and the body size of the Hokkaido and Sakhalin populations is similar to that of the Arctic warbler. In summary, the three species show minor morphological differences: the Arctic warbler is smaller and has grayish plumage, the Japanese leaf warbler is larger and has yellowish plumage, and the body size and plumage coloration of the Kamchatka leaf warbler is intermediate between those of the Arctic warbler and Japanese leaf warbler. As morphological differences are minor and overlap across species, no reliable method exists to distinguish the three species by appearance.

Molecular markers provide an alternative method for species identification. Mitochondrial DNA (mtDNA) sequences are one example of such markers. Because of their maternal inheritance (Dawid and Blackler 1972), small genome size (Moritz et al. 1987), and high mutation rate (Brown et al. 1979), mtDNA sequences can reveal genetic differentiation between species (Castro et al. 1998); therefore, they are widely used in phylogenetic studies. DNA barcoding is a global tool for specimen identification. Sequencing a fragment of the mtDNA gene and comparing the sequence with known sequences from a database can enable the accurate identification of species (Hebert et al. 2003). In birds, the mitochondrial cytochrome *c* oxidase subunit 1 (CO1) gene is used for DNA barcoding. The size of the CO1 gene used for DNA barcoding is approximately 648 bp. It is widely used for avian species identification and taxonomy research because of its small size, high success rate of sequencing, and low sequencing cost in addition to the availability of universal primer pairs (Hebert et al. 2004; Dove et al. 2008; Lijtmaer et al. 2011; Tavares et al. 2011; Colihueque et al. 2021). The Barcode of Life Data Systems (BOLD, <https://>

www.boldsystems.org/) is an online database hosting the DNA barcode data. Currently, more than 6000 bird species have sequences in this database. However, as taxonomic status changes through more research, some species classifications in the database may need to be updated (van den Burg and Vieites 2023).

During wintering, birds of the Arctic warbler species complex rarely sing; therefore, species identification by vocalization is not feasible for wintering birds. The plumage coloration of wintering birds is generally less bright. The occurrence of feather wear during migration further makes plumage coloration less species distinctive. In this study, we analyzed Arctic warbler species complex samples collected from a bird banding program. We used mitochondrial CO1 gene sequences derived from blood samples for species identification. For the first time, we confirmed that all three species visited Taiwan. Morphological measurements and photographs of the samples revealed that neither body size nor plumage coloration are reliable traits for distinguishing the species of the Arctic warbler species complex. In addition, we reanalyzed the CO1 gene sequences of the curated Arctic warbler species complex and found that many of their taxonomic statuses need to be updated.

## MATERIALS AND METHODS

### Bird banding and sample collection

Samples were collected from a bird banding program in Taroko National Taiwan, eastern Taiwan, that has been conducted since 2009. The environments are mainly grassland along the forest edge. We set up mist nets to catch birds. After capture, each bird was first fitted with a numbered metal ring on the tarsus. We then measured the weight, maximum wing length, tail length, bill length, and tarsus length of the birds. We collected approximately 20 µl blood by venipuncture from the brachial vein, and the blood samples were stored in absolute alcohol. We took pictures of each bird against a gray background, which was later used as a color reference to correct for white balance of the pictures. All the procedures could be completed in eight min. Subsequently, we immediately released the birds at the same site. Information on the samples, including sampling dates, locations, photographs, and sex, is provided in table S1 and the Barcode of Life Data System (ID: NDHUP001-23 to NDHUP036-23). Bird banding and sample collection were approved by Taroko National Park, Hualien Forest District Office, Forestry Bureau (now Hualien Branch, Forestry and Nature Conservation Agency, Department of Agriculture) and

the Hualien County Government.

### Downloading CO1 gene sequences of the Arctic warbler species complex from the database

We searched the CO1 gene sequences of the Arctic warbler species complex in the BOLD database by using *Phylloscopus borealis*, *Phylloscopus xanthodryans*, and *Phylloscopus examinandus* as keywords. We downloaded a total of 73 sequences from the database. After removing six sequences of poor quality and one sequence that was possibly a repeat sequence from the same individual, we obtained 66 sequences. Among these, 3 sequences were from the Japanese leaf warbler, 13 sequences were from the Kamchatka leaf warbler, and 50 sequences were from the Arctic warbler (Table S2). We identified collection sites and sampling dates from the database or from the original references. Before the Arctic warbler species complex is categorized into three species, all the birds are classified as the same species (*P. borealis*); therefore, some early submitted sequences of *P. borealis* may be derived from different species. In this study, to avoid potential mistakes in species identification, we selected 21 sequences that were collected during the breeding season (June to August, Billerman et al. 2022) and noted the collection sites. We then identified the species of these samples according to the breeding/collection sites. We identified a total of 12, 4, and 5 sequences in Arctic warbler, Japanese leaf warbler, and Kamchatka leaf warbler samples, respectively.

### DNA extraction, amplification, and sequencing

We used the FavorPrep™ Blood/Cultured Cell Genomic DNA Extraction Mini Kit (Favorgen, Taiwan) to extract genomic DNA from the blood samples of the Arctic warblers according to the manufacturer's instructions. We then performed polymerase chain reaction (PCR) to amplify the DNA fragments of sex chromosomes and the mitochondrial CO1 gene by using the extracted DNA as a template. For molecular sexing, we amplified a fragment of the chromodomain helicase DNA-binding protein (CHD) gene from sex chromosomes by using the primer pair of 2550F (5'-GTTACTGATTCGTCTACGAGA-3') and 2718R (5'-ATTGAAATGATCCAGTGCTTG-3') (Fridolfsson and Ellegren 1999). The reaction volume was 10 µl; it consisted of 0.5 µl DNA, 0.6 µl dNTP (2.5 mM of each nucleotide), 0.4 µl 2550F primer (10 mM), 0.4 µl 2718R primer (10 mM), 1.0 µl 10 × PCR buffer, 7.02 µl ddH<sub>2</sub>O and 0.08 Taq DNA polymerase (5 U/µl, Protech, Taiwan). The PCR program was as follows: 3 min at 95°C; 35 cycles of 30 s at 95°C, 40 s at 46°C, and 50 s

at 72°C; and a final extension step at 72°C for 5 min. We dyed the PCR products using EZ-Vision® Two, DNA Dye-as-Loading Buffer 6× (Protech, Taiwan) and then performed electrophoresis on 1.5% agarose gel (100 V, 30 min) using a Bio-100™ DNA Ladder (Protech) as a size reference. We examined the gels under ultraviolet light. Moreover, we identified the sex of the birds based on the presence or absence of the female-specific DNA fragment of the CHD-W gene.

We used the primer pair of AWCF1 (5'-CGCY TWAACAYTCYGCCATCTTACC-3') and AWC3 (5'-ATGCTCGGGTGTCTACGTCTAT-3') developed by Patel et al. (2010) to amplify a fragment of the mitochondrial CO1 gene sequence. The reaction volume was 20 µl; it consisted of 1.0 µl template DNA, 1.3 µl dNTP (Ex Taq™ dNTP, 2.5 mM, Takara), 0.6 µl AWCF1 primer (10 mM), 0.6 µl AWC3 primer (10 mM), 2.0 µl 10X PCR buffer (Ex Taq™ buffer, 10X, Takara, Japan), 1.3 µl MgCl<sub>2</sub> (25 mM), 12.9 µl ddH<sub>2</sub>O, and 0.3 µl Taq DNA polymerase (Takara Ex Taq™ polymerase, 5 U/µl, Takara, Japan). The PCR program was as follows: 3 min at 94°C; 35 cycles of 30 s at 94°C, 40 s at 63°C, 60 s at 72°C; and a final extension step at 72°C for 5 min. The PCR products were sequenced at the Cancer Progression Research Center, National Yang Ming Chiao Tung University. The sequences were aligned and edited using Sequencher 5.4.5 software (Gene Codes Corporation, Ann Arbor, Michigan, USA).

## CO1 gene sequence analyses

All sequences, including those downloaded from BOLD and those obtained from samples collected in this study, were aligned using Sequencher 5.4.5 software. DnaSP version 6.1203 (Rozas et al. 2017) was used to identify haplotypes and to calculate haplotype diversity ( $H_d$ ). Network 10 (Fluxus Engineering) was used to construct a haplotype network by using a medium-joining network algorithm (Bandelt et al. 1999).

## Species identification by plumage coloration

The plumage coloration of the banded Arctic warbler species complex was examined using photographs taken during the banding process. The photographs were first corrected for white balance, using PhotoDemon software (version 9.0). For some pictures without a gray background, we used the same software to automatically correct white balance. Colors of upper regions (head, back, and wing) and underparts (throat and breast) were used to identify the species. For each bird, the upper regions were denoted as having the coloration of olive green, grayish olive green, and light olive green by using the photographs provided by Round et al. (2016) as reference. The underparts were denoted as having the coloration of white, yellow, and light yellow. Figure 1 illustrates some plumage coloration. Species identification was performed based



**Fig. 1.** The plumage coloration of the Arctic Warbler species complex, showing olive green upperparts (A), grayish green upperparts (B), light olive-green upperparts (C), paler and whitish underparts (D), variably yellowish throat and underparts (E), and bright yellow throat and underparts (F).

on coloration (del Hoyo et al. 2020a b; Lowther and Sharbaugh 2020; Hsiao and Li 2017). All coloration assignment and species identification procedures were performed by SJC. The results were compared with those of genetic analyses.

### Morphological measurements

According to the results of molecular sexing and CO1 species identification, we explored sexual dimorphism in the measurements of Arctic warblers. Two samples had missing values for some traits; we therefore excluded them from the analysis. We performed principal component analysis (PCA) to assess the differences in the size of the samples.

## RESULTS

A total of 36 birds from the Arctic warbler species complex were banded. Molecular sexing revealed 17 male birds and 19 female birds.

### Species identification by CO1 gene sequences


In this study, 607 bp fragments of CO1 gene sequences were obtained from 102 samples, including

66 sequences downloaded from BOLD and 36 sequences obtained from Taiwanese birds. The sequence set contained 49 variable sites, of which 44 were located at the third nucleotide of a codon. In total, 27 haplotypes were identified.

Based on the 21 CO1 gene sequences of the Arctic warbler species complex from BOLD, which were collected from the breeding area during the breeding season, three haplogroups were identified. The haplogroups matched well with the sampling sites (Fig. 1). This indicates that CO1 gene sequences are an ideal marker for distinguishing Arctic warblers, Japanese leaf warblers, and Kamchatka leaf warblers.

Using the haplotype network, we identified the species of the 36 samples collected in this study. Specifically, thirty-two of them were Arctic warblers (14 males and 18 females), two were Japanese leaf warblers (both males), and two were Kamchatka leaf warblers (one male and one female) (Fig. 2, Table S1).

Reassessment of the CO1 gene sequences of the Arctic Warbler species complex deposited in BOLD revealed that the species names of many samples need to be updated. Three Japanese leaf warbler samples in BOLD should be reclassified as Arctic warblers. Of the 50 Arctic warbler samples in BOLD, five should be reclassified as Japanese leaf warblers, and 13 should be reclassified as Kamchatka leaf warblers. The 13



**Fig. 2.** Haplotype networks of the Arctic warbler species complex (*Phylloscopus borealis* spp.) constructed using fragments of mitochondrial CO1 gene sequences (607 bp). Blue indicates samples from BOLD, which were collected from breeding areas during the breeding season. Yellow represents the Taiwanese samples obtained from this study. The size of the circle is proportional to the sample size.



Kamchatka leaf warblers in BOLD are all correctly classified. In total, the taxonomic status of 31.8% of the samples (21/66) needs to be updated (Table S2).

On combining sequences from the BOLD database and the Taiwanese samples, sixteen of 27 haplotypes were identified from Arctic warblers ( $n = 67$ ,  $H_d = 0.767$ ). Moreover, four haplotypes were identified from Japanese leaf warblers ( $n = 7$ ,  $H_d = 0.81$ ), and seven were identified from Kamchatka leaf warblers ( $n = 28$ ,  $H_d = 0.743$ ) (Tables S1, S2).

Species identification by plumage coloration and size

Using the results of genetic species identification, we examined the plumage coloration and morphological measurement data of the three species. Both traits showed extensive overlap among species; therefore, they cannot be used for species identification. Of the 30 samples for which photo records were available, the species of only 50% (15/30) were correctly identified by plumage coloration (Table 1, Table S2). Identification errors occurred for all three species.

We noted sexual dimorphism in the morphological measurements of Arctic warblers. Male birds were significantly larger than female birds in all measurements, except for the bill length (Table 2). Because only two Japanese leaf warblers (both male

birds) and two Kamchatka leaf warblers (one male and one female) were identified, we could not compare sexual dimorphism for these two species. In PCA, both PC1 and PC2 explained 71.8% of the variation (44.9% and 26.9% for PC1 and PC2, respectively). PCA of all samples revealed that the measurements of the three species greatly overlapped. However, Kamchatka leaf warblers tended to be larger in both PC1 and PC2, whereas Japanese leaf warblers tended to be smaller in PC2 (Fig. 3).




Fig. 3. Principal component analysis of measurement data of the Arctic warbler species complex. Samples marked with “J” are Japanese leaf warblers, those marked with “K” are Kamchatka leaf warblers, and the rest are Arctic warblers.

Table 1. Comparison of species identification by mitochondrial CO1 gene sequences and plumage coloration. Numbers in bold indicate the number of birds that were identified as the same species via both methods

Species identified by CO1 gene sequences	N	Species identified by plumage coloration		
		Arctic warbler	Japanese leaf warbler	Kamchatka leaf warbler
Arctic warbler	26	<b>13</b>	2	11
Japanese leaf warbler	2	1	<b>1</b>	0
Kamchatka leaf warbler	2	1	0	<b>1</b>

Table 2. Morphological measurements of the Arctic warbler species complex samples collected from Taiwan

Measurements	Species					
	Arctic warbler			Kamchatka leaf warbler		Japanese leaf warbler
	Male (n = 14)	Female (n = 16)	t test	Male (n = 1)	Female (n = 1)	Male (n = 2)
Weight (g)	9.67 ± 1.09	8.93 ± 1.10	t = 1.85, p = 0.04	10.4	11.8	9.15 ± 0.78
Maximum wing length (mm)	66.11 ± 2.65	62.56 ± 2.57	t = 3.70, p < 0.001	69.0	66.0	67.50 ± 0.71
Tail length (mm)	47.46 ± 3.23	44.63 ± 3.21	t = 2.41, p = 0.012	44.5	48.0	51.25 ± 0.35
Bill length (mm)	9.43 ± 0.46	9.29 ± 0.41	t = 0.91, p = 0.19	9.67	10.19	8.65 ± 0.29
Tarsus length (mm)	19.32 ± 0.53	18.78 ± 0.63	t = 2.57, p = 0.008	20.35	19.71	19.61 ± 0.89

## DISCUSSION

Taiwan is located in the middle of the East Asian-Australasian Avian Flyway. The annual latitudinal migration of nearly 700 species of birds occurs along this flyway (Kirby et al. 2008). According to the latest checklist, no confirmed record of the Japanese leaf warbler or Kamchatka leaf warbler currently in Taiwan (Ding et al. 2023). In this study, we successfully identified the three species of the Arctic warbler species complex based on mitochondrial CO1 gene sequences. Our results present strong evidence of their presence in Taiwan.

All our Japanese leaf warbler and Kamchatka leaf warbler samples were banded during the migration season: three were collected in October and one was collected in May. Given their low number and appearance in the migration season, both species appear to be rare transients at our study site (eastern Taiwan). More samples from other regions of Taiwan would help reveal their population status in Taiwan.

During nonbreeding season, the Arctic warbler species complex cannot be distinguished by species-specific songs. Although several studies have reported differences in plumage coloration (del Hoyo et al. 2020a b; Lowther and Sharbaugh 2020), we failed to distinguish the three species based on the plumage coloration of our samples. A similar result was reported by Round et al. (2016); they found that the plumage coloration of some wintering Arctic warblers was very similar to that of the Kamchatka leaf warbler. For birds in migration, feather abrasion, changes in food composition, and different molting stages may make species identification by morphology more difficult.

According to the results of PCA, we did not find considerable disparity in body size among the three species. The Kamchatka leaf warbler tended to be larger than the Arctic warbler and Japanese leaf warbler. This does not fully conform to the findings of previous studies (Saitoh et al 2008; Round et al. 2015 2016), which have reported that the body size of the Kamchatka leaf warbler is intermediate of the size of the Arctic warbler and Japanese leaf warbler. The body weight of migratory birds may fluctuate greatly, owing to fat deposition and depletion along the migratory routes. Therefore, there may be high variation in weight either within or among species. In addition, we noted significant sexual dimorphism in the body size of Arctic warblers. However, we did not have sufficient samples to test whether such dimorphism exists in the other two species. If all three species exhibit sexual dimorphism in their body size, species identification by body size will become more unreliable.

For closely related species, incomplete lineage

sorting and introgression through secondary contact may lead to the three species not necessarily presenting reciprocal monophyly. Given the fact that the three species are not sympatric during breeding season, and they have distinct species-specific vocalizations, genetic introgression through hybridization seems less likely. In addition, in a study on the phylogeny of Phylloscopidae warblers, Alström et al. (2018) did not find evidence of incomplete lineage sorting in the mitochondrial cytochrome *b* gene. However, they found evidence of incomplete lineage sorting in nuclear genes in some lineages. In our samples, the three species cannot be distinguished by morphology, and therefore we cannot rule out the possibility of incomplete lineage sorting among them.

In addition to examining the Taiwanese samples, we reassessed the sequences deposited in BOLD, and the results revealed that a high proportion (31.8%) of the samples need to revise their species names. Nearly all the Japanese leaf warblers and Kamchatka leaf warblers misidentified as Arctic warblers in BOLD were submitted before the splitting of the Arctic warbler species complex into the Arctic warbler, Japanese leaf warbler, and Kamchatka leaf warbler (Clements et al. 2014). Three Arctic warblers have been misidentified as Japanese leaf warblers in BOLD. These samples were collected from South Korea (Park et al. 2011), where the Arctic warbler species complex is regarded as transient (it does not breed or winter there) (del Hoyo et al. 2020a b; Lowther and Sharbaugh 2020). These samples may be identified by morphology, which was proven to be ambiguous for species identification in this study.

A biological database serves as an invaluable resource for biodiversity research. However, as the amount of data increases, outdated taxonomic statuses are a potential problem. For example, in a recent study, van den Burg et al. (2020) found that approximately 13% of amphibian cytochrome *b* records in GenBank are taxonomically outdated. Päckert (2022) found that some Charadriidae CO1 gene sequences in GenBank may not correspond to their taxon names, which may lead to inaccurate classification results. We propose that while extracting data from BOLD, taxonomic updates should be considered.

## CONCLUSIONS

In this study, we confirmed that the CO1 gene sequence can effectively identify the three species of the Arctic warbler species complex. Applying this method to samples collected from eastern Taiwan, for the first time, we confirmed the presence of the Japanese leaf warbler and Kamchatka leaf warbler in Taiwan,

adding two new species to the country's avifauna. In addition, our results highlighted that the morphological (both measurement data and plumage coloration) differences among species, as described in the literature and guidebooks, are not reliable for distinguishing the species during the migration/wintering season. Moreover, as the taxonomic status may change over time, caution should be taken when extracting sequence data from a database (such as BOLD).

**Acknowledgments:** The long-term bird banding project was funded by Taroko National Park (TNP) to YCH. The sequence analyses were funded by a student research grant of TNP to SJC. We acknowledge all members of Bird Lab, National Dong Hwa University, for assisting with field work. We thank Pei-Chun Shih for preparing the figures.

**Authors' contributions:** Both authors contributed equally to field work, data analyses, and manuscript writing. SJC performed morphological measurements and DNA sequencing.

**Competing interests:** The authors declare that they have no competing interests.

**Availability of data and materials:** Sequence data are submitted to BOLD. The metadata of the samples are attached in the Supplementary Materials.

**Consent for publication:** All the authors consent to the publication of this manuscript.

**Ethics approval consent to participate:** Bird banding and sample collection were approved by Taroko National Park, Hualien Forest District Office, Forestry Bureau (now Hualien Branch, Forestry and Nature Conservation Agency, Department of Agriculture), and the Hualien County Government.

## REFERENCES

- Adams EM, Stenhouse IJ, Gilbert AT, Boelsma J, Gress G, Weidensaul CS, Grigsby C, Williams EJ, Phillips L, McIntyre CL. 2022. The first documentation of the Nearctic-Paleotropical migratory route of the Arctic Warbler. *Ecol Evol* **12**:e09223. doi:10.1002/ece3.9223.
- Alström P, Saitoh T, Williams D, Nishiumi I, Shigeta Y, Ueda K, Irestedt M, Björklund M, Olsson U. 2011. The Arctic Warbler *Phylloscopus borealis* – three anciently separated cryptic species revealed. *Ibis* **153**:395–410. doi:10.1111/j.1474-919X.2011.01116.x.
- Alström P, Rheindt FE, Zhang R, Zhao M, Wang J, Zhu X, Gwee CY, Hao Y, Ohlson J, Jia C, Prawiradilaga DM, Ericson PGP, Lei F, Olsson U. 2018. Complete species-level phylogeny of the leaf warbler (Aves: Phylloscopidae) radiation. *Mol Phylogenet Evol* **126**:141–152. doi:10.1016/j.ympev.2018.03.031.
- Bandelt HJ, Forster P, Röhl A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol* **16**:37–48. doi:10.1093/oxfordjournals.molbev.a026036.
- Billerman SM, Keeney BK, Rodewald PG, Schulenberg TS (eds). 2022. *Birds of the World*. Cornell Laboratory of Ornithology, Ithaca, New York. Available at: <https://birdsoftheworld.org/bow/home>. Accessed 6 Oct. 2023.
- Brown WM, George M Jr, Wilson AC. 1979. Rapid evolution of animal mitochondrial DNA. *Proc Natl Acad Sci USA* **76**:1967–1971. doi:10.1073/pnas.76.4.1967.
- Castro JA, Picornell A, Ramon M. 1998. Mitochondrial DNA: a tool for population genetics studies. *Int Microbiol* **1**:327–332.
- Cazalis V, Princé K, Minhoub J-B, Kelly J, Butchart SHM, Rodrigues ASL. 2020. Effectiveness of protected areas in conserving tropical forest birds. *Nat Commun* **11**:4461. doi:10.1038/s41467-020-18230-0.
- Clements JF, Schulenberg TS, Iliff MJ, Roberson D, Fredericks TA, Sullivan BL, Wood CL. 2014. *The eBird/Clements Checklist of Birds of the World: Version 6.9*.
- Colihueque N, Gantz A, Parraquez M. 2021. Revealing the biodiversity of Chilean birds through the *COI* barcode approach. *ZooKeys* **1016**:143–161. doi:10.3897/zookeys.1016.51866.
- Dawid IB, Blackler AW. 1972. Maternal and cytoplasmic inheritance of mitochondrial DNA in *Xenopus*. *Dev Biol* **29**:152–161. doi:10.1016/0012-1606(72)90052-8.
- del Hoyo J, Collar N, Christie DA. 2020a. Japanese Leaf Warbler (*Phylloscopus xanthodryas*), version 1.0. In: del Hoyo J, Elliott A, Sargatal J, Christie DA, de Juana E (eds) *Birds of the World*. Cornell Lab of Ornithology, New York. doi:10.2173/bow.arcwar3.01.
- del Hoyo J, Collar N, Christie DA. 2020b. Kamchatka Leaf Warbler (*Phylloscopus examinandus*), version 1.0. In: del Hoyo J, Elliott A, Sargatal J, Christie DA, de Juana E (eds) *Birds of the World*. Cornell Lab of Ornithology, New York. doi:10.2173/bow.arcwar2.01.
- Ding T-S, Juan C-S, Lin R-S, Tsai Y-J, Wu J-L, Wu J, Yang Y-H. 2023. The 2023 CWF Checklist of the Birds of Taiwan. Taiwan Wild Bird Federation, Taipei.
- Dove CJ, Rotzel NC, Heacker M, Weigt LA. 2008. Using DNA barcodes to identify bird species involved in birdstrikes. *J Wildlife Manage* **72**:1231–1236. doi:10.2193/2007-272.
- Exantus J-M, Cézilly F. 2023. Composition of avian assemblage in a protected forested area in Haiti: Evidence for recent decline of both forest-dependent and insectivore species. *Glob Ecol Conserv* **46**:e02607. doi:10.1016/j.gecco.2023.e02607.
- Fridolfsson A-K, Ellegren H. 1999. A simple and universal method for molecular sexing of non-ratite birds. *J Avian Biol* **30**:116–121. doi:10.2307/3677252.
- Hebert PDN, Ratnasingham S, deWaard JR. 2003. Barcoding animal life: cytochrome *c* oxidase subunit 1 divergences among closely related species. *Proc Biol Sci* **270**:S96–S99. doi:10.1098/rsbl.2003.0025.
- Hebert PDN, Stoeckle MY, Zemlak TS, Francis CM. 2004. Identification of birds through DNA barcodes. *PLoS Biol* **2**:e312. doi:10.1371/journal.pbio.0020312.
- Hsiao M-C, Li C-L. 2017. *A Field Guide to the Birds of Taiwan*. Wild Bird Society of Taipei, Taipei, Taiwan.
- Kirby JS, Stattersfield AJ, Butchart SHM, Evans MI, Grimmett RFA, Jones VR, O'Sullivan J, Tucker GM, Newton I. 2008. Key conservation issues for migratory land- and waterbird species on the world's major flyways. *Bird Conserv Int* **18**:S49–S73. doi:10.1017/S0959270908000439.
- Lijtmaer DA, Kerr KCR, Barreira AS, Hebert PDN, Tubaro PL. 2011.



- DNA barcode libraries provide insight into continental patterns of avian diversification. PLoS ONE **6**:e20744. doi:10.1371/journal.pone.0020744.
- Lowther PE, Sharbaugh S. 2020. Arctic Warbler (*Phylloscopus borealis*), version 1.0. In: Billerman SM (ed) Birds of the World. Cornell Lab of Ornithology, New York, USA. doi:10.2173/bow.arcwarl.01.
- Mace GM. 2004. The role of taxonomy in species conservation. Philos Trans R Soc Lond B Biol Sci **359**:711–719. doi:10.1098/rstb.2003.1454.
- Martens J. 2010. A preliminary review of the leaf warbler genera *Phylloscopus* and *Seicercus*. Br Ornithol Club Occas Publ **5**:41–116.
- Moritz C, Dowling TE, Brown WM. 1987. Evolution of animal mitochondrial DNA: Relevance for population biology and systematics. Annu Rev Ecol Syst **18**:269–292. doi:10.1146/annurev.es.18.110187.001413.
- Patel S, Waugh J, Millar CD, Lambert DM. 2010. Conserved primers for DNA barcoding historical and modern samples from New Zealand and Antarctic birds. Mol Ecol Resour **10**:431–438. doi:10.1111/j.1755-0998.2009.02793.x.
- Päckert M. 2022. Free access of published DNA sequences facilitates regular control of (meta-) data quality – an example from shorebird mitogenomes (Aves, Charadriiformes: *Charadrius*). Ibis **164**:336–342. doi:10.1111/ibi.13005.
- Park H-Y, Yoo HS, Jung G, Kim C-B. 2011. New DNA barcodes for identification of Korean birds. Genes Genomics **33**:91–95. doi:10.1007/s13258-010-0089-3.
- Reeves AB, Drovetski SV, Fadeev IV. 2008. Mitochondrial DNA data imply a stepping-stone colonization of Beringia by arctic warbler *Phylloscopus borealis*. J Avian Biol **39**:567–575. doi:10.1111/j.0908-8857.2008.04421.x.
- Rozas J, Ferrer-Mata A, Sanchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, Sanchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism analysis of large data sets. Mol Biol Evol **34**:3299–3302. doi:10.1093/molbev/msx248.
- Round PD, Pierce AJ, Nualsri C, Sankamethawee W. 2015. A record of Japanese Leaf Warbler *Phylloscopus xanthodryas* in Thailand. BirdingASIA **23**:133–134.
- Round PD, Pierce AJ, Saitoh T, Shigeta Y. 2016. Addition of Kamchatka Leaf Warbler *Phylloscopus examinandus* and Sakhalin Leaf Warbler *P. borealoides* to Thailand's avifauna. Bull Jpn Bird Banding Assoc **28**:9–21. doi:10.14491/jbba.MS076.
- Rutt CL, Jirinec V, Cohn-Haft M, Laurance WF, Stouffer PC. 2019. Avian ecological succession in the Amazon: A long-term case study following experimental deforestation. Ecol Evol **9**:13850–13861. doi:10.1002/ece3.5822.
- Saitoh T, Shigeta Y, Ueda K. 2008. Morphological differences among populations of the Arctic Warbler with some intraspecific taxonomic notes. Ornithol Sci **7**:135–142. doi:10.2326/1347-0558-7.2.135.
- Saitoh T, Alström P, Nishiumi I, Shigeta Y, Williams D, Olsson U, Ueda K. 2010. Old divergences in a boreal bird supports long-term survival through the Ice Ages. BMC Evol Biol **10**:35. doi:10.1186/1471-2148-10-35.
- Severinghaus LL, Ding TS, Fang WH, Ling WH, Tsai MC, Yen CW. 2012. The Avifauna of Taiwan, 2nd edn, vol 3. Forest Bureau, Council of Agriculture, Taipei, Taiwan.
- Tavares ES, Gonçalves P, Miyaki CY, Baker AJ. 2011. DNA barcode detects high genetic structure within Neotropical bird species. PLoS ONE **6**:e28543. doi:10.1371/journal.pone.0028543.
- van den Burg MP, Vieites DR. 2023. Bird genetic databases need improved curation and error reporting to NCBI. Ibis **165**:472–481. doi:10.1111/ibi.13143.
- van den Burg MP, Herrando-Pérez S, Vieites DR. 2020. ACDA, a global database of amphibian cytochrome-*b* sequences using reproducible curation for GenBank records. Sci Data **7**:268. doi:10.1038/s41597-020-00598-9.
- Vaurie C. 1954. Systematic Notes on Palearctic Birds. No. 9 Sylviinae: the Genus *Phylloscopus*. Am Mus Novit **1685**:1–24.
- Withrow JJ, Gibson DD, Gerasimov Y, Gerasimov N, Shestopalov A, Winker K. 2016. Occurrence and taxonomy of Arctic Warblers (*Phylloscopus borealis*) *sensu lato* in North America. Wilson J Ornithol **128**:268–277. doi:10.1676/wils-128-02-268-277.1.

## Supplementary materials

**Table S1.** Information about the Arctic warbler species complex samples collected in Taiwan that were used in this study. (download)

**Table S2.** Information about the Arctic warbler species complex samples downloaded from the Barcode of Life Data System. (download)